


#2

REC'D 22 SEP 2003

WIPO

PCT

P1 1065078



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

September 16, 2003

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/400,352

FILING DATE: July 31, 2002

RELATED PCT APPLICATION NUMBER: PCT/US03/23821



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

M. Sias

M. SIAS
Certifying Officer

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

08-02-29 0035 073102 A/Prov.

07/31/02
 1-800 U.S. PTO

Please type a plus sign (+) inside this box



PTO/SB/16 (10-01)

Approved for use through 10/31/2002. OMB 0851-0032

U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

Express Mail Label No. EU465991597US

1-800 U.S. PTO
 60/400352
 07/31/02

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
ERIC DEGUYON		TAYLOR		NEWARK, DELAWARE	
<input type="checkbox"/> Additional inventors are being named on the _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
METHOD FOR PREPARING FUSED OXAZINONES					
Direct all correspondence to:			CORRESPONDENCE ADDRESS		
<input checked="" type="checkbox"/> Customer Number OR			23906 Type Customer Number Here		
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages 73		<input type="checkbox"/> CD(s), Number _____		<input type="checkbox"/> Other (specify) Return Receipt Postcard	
<input type="checkbox"/> Drawing(s) Number of Sheets _____		<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76			
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.					
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees					
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number:				04-1928	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				160.00	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

Respectfully submitted,

Date 07/31/02

SIGNATURE

TYPED or PRINTED NAME DAVID E. HEISER

TELEPHONE 302-892-1926

REGISTRATION NO. 31,366
(If appropriate)

Docket Number

BA9308 US PRV

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C., 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

**FEE TRANSMITTAL
for FY 2002**

Patent fees are subject to annual revision

☐ Applicant Claims small entity status. See 37 CFR 1.27**TOTAL AMOUNT OF PAYMENT (\$)** 160**Complete If Known**

Application Number	UNKNOWN
Filing Date	CONCURRENTLY HEREWITH
First Named Inventor	ERIC DEGUYON TAYLOR
Examiner Name	UNKNOWN
Group / Art Unit	UNKNOWN
Attorney Docket No.	BA9308 US PRV

METHOD OF PAYMENT (check all that apply)☐ Check ☐ Credit card ☐ Money Order ☐ Other ☐ None☒ Deposit AccountDeposit
Account
Number

04-1928

Deposit
Account
Name

E. I. du Pont de Nemours and Company

The Commissioner is authorized to: (check all that apply)

☒ Charge fee(s) indicated below ☒ Credit any overpayments
☒ Charge any additional fee(s) during the pendency of this application
☐ Charge fee(s) indicated below, except for the filing fee to the above-identified deposit account
FEE CALCULATION**1. BASIC FILING FEE**

Large Entity		Small Entity		Fee Description	Fee Paid
Fee Code	Fee (\$)	Fee Code	Fee (\$)		
101	740	201	370	Utility filing fee	
108	330	208	165	Design filing fee	
107	510	207	255	Plant filing fee	
108	740	208	370	Reissue filing fee	
114	160	214	80	Provisional filing fee	160
SUBTOTAL (1)					(\$) 160

2. EXTRA CLAIM FEES

Total Claims		-20	=	0	X	18	=	0
Independent Claims		-3	=	0	X	84	=	0
Multiple Dependent	<input type="checkbox"/>				X	280	=	0

Large Entity		Small Entity		Fee Description	Fee Paid
Fee Code	Fee (\$)	Fee Code	Fee (\$)		
103	18	203	9	Claims in excess of 20	
102	84	202	42	Independent claims in excess of 3	
104	280	204	140	Multiple dependent claim, if not paid	
109	84	209	42	** Reissue independent claims over original patent	
110	18	210	9	** Reissue claims in excess of 20 and over original patent	

SUBTOTAL (2)

(\$) 0

**or number previously paid, if greater; For Reissues, see above

FEE CALCULATION (continued)**3. ADDITIONAL FEES**

Large Entity		Small Entity		Fee Description	Fee Paid
Fee Code	Fee (\$)	Fee Code	Fee (\$)		
105	130	205	65	Surcharge - late filing fee or oath	
127	50	227	25	Surcharge - late provisional filing fee or cover sheet	
139	130	139	130	Non-English specification	
147	2,520	147	2,520	For filing a request for reexamination	
112	920*	112	920*	Requesting publication of SIR prior to Examiner action	
113	1,840*	113	1,840*	Requesting publication of SIR after Examiner action	
115	110	215	55	Extension for reply within first month	
116	400	216	200	Extension for reply within second month	
117	920	217	460	Extension for reply within third month	
118	1,440	218	720	Extension for reply within fourth month	
128	1,960	228	980	Extension for reply within fifth month	
119	320	219	160	Notice of Appeal	
120	320	220	160	Filing a brief in support of an appeal	
121	280	221	140	Request for oral hearing	
138	1,510	138	1,510	Petition to institute a public use proceeding	
140	110	240	55	Petition to revive - unavoidable	
141	1,280	241	640	Petition to revive - unintentional	
142	1,280	242	640	Utility issue fee (or reissue)	
143	480	243	230	Design issue fee	
144	620	244	310	Plant issue fee	
122	130	122	130	Petitions to the Commissioner	
123	50	123	50	Processing fee under 37 CFR 1.17(q)	
128	180	128	180	Submission of Information Disclosure Stmt	
581	40	581	40	Recording each patent assignment per property (times number of properties)	
146	740	246	370	Filing a submission after final rejection (37 CFR § 1.129(a))	
149	740	249	370	For each additional invention to be examined (37 CFR § 1.129(b))	
179	740	279	370	Request for Continued Examination (RCE)	
169	900	169	900	Request for expedited examination of a design application	

Other fee (specify) _____

*Reduced by Basic Filing Fee Paid

SUBTOTAL (3)

(\$) 0

SUBMITTED BY**Complete (if applicable)**

Name (Print/Type)	DAVID E HEISER	Registration No. Attorney/Agent	31,366	Telephone	302-892-1926
Signature	<i>David E Heiser</i>			Date	JULY 31, 2002

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

TITLE

METHOD FOR PREPARING FUSED OXAZINONES

BACKGROUND OF THE INVENTION

5 A need exists for additional methods for preparing fused oxazinones. Such compounds include intermediates for the preparation of crop protection agents, pharmaceuticals and other fine chemicals.

Fused oxazinones have been prepared by a variety of methods. For example, *N*-acylanthranilic acids have been treated with acetic anhydride, anthranilic acids have been treated with carboxylic acid anhydrides, and anthranilic acids and carboxylic acids have been
10 coupled in the presence of various dehydrating agents (see G. M. Coppola, *J. Heterocyclic Chemistry* 1999, 36, 563-588). Fused oxazinones have also been prepared by treatment of *ortho*-amino carboxylic acids with carboxylic acid chlorides in the presence of base (see e.g. Jakobsen et al, *Biorganic and Medicinal Chemistry*, 2000, 8, 2803-2812 and Jakobsen et al., *Biorganic and Medicinal Chemistry* 2000, 8, 2095-2103). Benzoxazinones have also been
15 prepared by treatment of a carboxylic acid with a sulfonyl chloride and then treatment with an anthranilic acid (see D. V. Ramana and E. Kantharaj, *Org. Prep. Proced. Int.*, 1993, 25, 588).

SUMMARY OF THE INVENTION


20 This invention provides a method for preparing a fused oxazinone. This method comprises:

(1) contacting a carboxylic acid with a sulfonyl chloride in the presence of an optionally substituted pyridine compound, the nominal mole ratio of sulfonyl chloride to carboxylic acid being from about 0.75 to 1.5;

25 (2) contacting the mixture prepared in (1) with an *ortho*-amino aromatic carboxylic acid in the presence of an optionally substituted pyridine compound, the nominal mole ratio of the *ortho*-amino aromatic carboxylic acid to carboxylic acid charged in (1) being from about 0.8 to 1.2; and


(3) adding additional sulfonyl chloride to the mixture prepared in (2), the nominal mole ratio of additional sulfonyl chloride added in (3) to carboxylic acid charged in (1) being
30 at least about 0.5.

This invention also relates to a method for preparing a compound of Formula III,




wherein

- X is N or CR⁶;
 Y is N or CH;
 R¹ is H;
 R² is H or CH₃;
 R³ is C₁-C₆ alkyl;
 R⁴ is C₁-C₄ alkyl or halogen;
 R⁵ is H, C₁-C₄ alkyl, C₁-C₄ haloalkyl or halogen;
 R⁶ and R⁷ are independently H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, halogen, CN or C₁-C₄ haloalkoxy;
 R⁸ is H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, C₃-C₆ cycloalkylamino, (C₁-C₄ alkyl)(C₃-C₆ cycloalkyl)amino, C₂-C₄ alkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₂-C₆ alkylaminocarbonyl, C₃-C₈ dialkylaminocarbonyl or C₃-C₆ trialkylsilyl; and
 R⁹ is CF₃, OCF₃, OCHF₂, OCH₂CF₃, S(O)_pCF₃, S(O)_pCHF₂ or halogen;
- using a compound of Formula 1a




This method is characterized by preparing the compound of Formula 1a (i.e. a subgenus of Formula 1 described below) by the method as indicated above. Formula 1a compounds may be prepared using a compound of the formula $LS(O)_2Cl$ (wherein L is selected from alkyl, haloalkyl and phenyl optionally substituted with from one to three substituents independently selected from alkyl or halogen) as the sulfonyl chloride, a compound of
 5 Formula 2' as the carboxylic acid, and a compound of Formula 5' as the *ortho*-amino aromatic carboxylic acid



DETAILED DESCRIPTION OF THE INVENTION

One aspect of this invention involves the use of optionally substituted pyridine compounds together with sulfonyl chloride compounds to provide advantageous production of fused oxazinones. The optionally substituted pyridine compound is used (a) to facilitate contact between the carboxylic acid and the sulfonyl chloride compound and (b) to facilitate contact of the resulting mixture with the *ortho*-amino carboxylic acid. The sulfonyl chloride
 15 compound is used both (a) as a reactant to facilitate coupling of the carboxylic acid with the *ortho*-amino carboxylic acid and (b) as a reactant to facilitate ring closure to form the fused oxazinone. These aspects provide effective production of the fused oxazinone while limiting the amounts of the carboxylic acid and the *ortho*-amino carboxylic acid that are consumed during the formation of the fused oxazinone. This can be especially important when the
 20 carboxylic acid and/or the *ortho*-amino carboxylic acid are valuable, complex and/or difficult to obtain. For example, this invention may be used for preparing a compound of Formula 1




wherein

25 J is an optionally substituted carbon moiety; and

4


K is, together with the two contiguous linking carbon atoms, a fused phenyl ring or a fused 5- or 6-membered heteroaromatic ring, each ring optionally substituted.

More particularly, a compound of Formula 1 may be prepared by a method comprising (1) contacting a carboxylic acid of Formula 2



5


with a sulfonyl chloride of Formula 4



wherein L is selected from alkyl, haloalkyl, and phenyl optionally substituted with from one to three substituents independently selected from alkyl or halogen;

10 in the presence of an optionally substituted pyridine compound;

(2) contacting the mixture prepared in (1) with an *ortho*-amino carboxylic acid of Formula 5



in the presence of an optionally substituted pyridine compound; and

(3) contacting the mixture prepared in (2) with additional sulfonyl chloride of Formula 4.

15 In the recitations herein, the term "carbon moiety" refers to a radical in which a carbon atom is connected to the backbone of the fused oxazinone ring of Formula 1 and the carboxylic acid group of Formula 2. As the carbon moiety L is separated from the reaction center, it can encompass a great variety of carbon-based moieties that can be prepared by modern methods of synthetic organic chemistry. The method of this invention is generally applicable to prepare a wide range of compounds of Formula 1.

20

"Carbon moiety" thus includes alkyl, alkenyl and alkynyl moieties, which can be straight-chain or branched.

25

"Carbon moiety" also includes carbocyclic and heterocyclic rings, which can be saturated, partially saturated, or completely unsaturated. The carbocyclic and heterocyclic rings of a carbon moiety group can form polycyclic ring systems comprising multiple rings connected together. The term "carbocyclic ring" denotes a ring wherein the atoms forming the ring backbone are selected only from carbon. "Saturated carbocyclic" refers to a ring having a backbone consisting of carbon atoms linked to one another by single bonds; unless otherwise specified, the remaining carbon valences are occupied by hydrogen atoms. The

term "hetero" in connection with rings or ring systems refers to a ring or ring system in which at least one ring atom is not carbon and which can contain from one to four heteroatoms independently selected from the group consisting of nitrogen, oxygen and sulfur, provided that each ring contains no more than 4 nitrogens, no more than 2 oxygens and no more than 2 sulfurs. When the carbon moiety consists of a heterocyclic ring or ring system, it is connected to the backbone of the fused oxazinone ring of Formula 1 and the carboxylic acid group of Formula 2 through any available carbon ring atom by replacement of a hydrogen on said carbon atom. Heterocyclic rings or ring systems may also be connected to the carbon moiety through any available carbon or nitrogen atom by replacement of a hydrogen on said carbon or nitrogen atom.

Although there is no definite limit to the size of Formula 1 suitable for the process of the invention, typically Formula 1 comprises 9-100, more commonly 9-50, and most commonly 9-25 carbon atoms, and 3-25, more commonly 3-15, and most commonly 3-10 heteroatoms. Heteroatoms are atoms other than carbon or hydrogen and are commonly selected from halogen, oxygen, sulfur, nitrogen, phosphorus and silicon. Three heteroatoms in Formula 1 are the nitrogen atom and the two oxygen atoms in the oxazinone moiety. When K is a fused heteroaromatic ring or when the carbon moiety comprises a heterocyclic ring, additional heteroatoms are contained therein. Substituents attached to the K ring or the carbon moiety may also contain additional heteroatoms.

Unsaturated rings can be aromatic if Hückel's rule is satisfied. "Aromatic" indicates that each of the ring atoms is essentially in the same plane and has a *p*-orbital perpendicular to the ring plane, and in which $(4n + 2) \pi$ electrons, when *n* is 0 or a positive integer, are associated with the ring to comply with Hückel's rule. The term "aromatic ring system" denotes fully unsaturated carbocycles and heterocycles in which at least one ring of a polycyclic ring system is aromatic. The term "aromatic carbocyclic ring or ring system" includes fully aromatic carbocycles and carbocycles in which at least one ring of a polycyclic ring system is aromatic (e.g., phenyl, naphthyl and 1,2,3,4-tetrahydronaphthalenyl). The term "nonaromatic carbocyclic ring or ring system" denotes fully saturated carbocycles as well as partially or fully unsaturated carbocycles where the Hückel rule is not satisfied by any of the rings in the ring system. The terms "heteroaromatic ring or ring system" and "aromatic fused heterobicyclic ring system" includes fully aromatic heterocycles and heterocycles in which at least one ring of a polycyclic ring system is aromatic (where aromatic indicates that the Hückel rule is satisfied). The term "nonaromatic heterocyclic ring or ring system" denotes fully saturated heterocycles as well as partially or fully unsaturated heterocycles where the Hückel rule is not satisfied by any of the rings in the ring system. The term "aryl" denotes a carbocyclic or heterocyclic ring or ring system in which at least one ring is aromatic, and the aromatic ring provides the connection to the remainder of the molecule.

The carbon moiety specified for J is optionally substituted. Also, the K ring moieties of Formulae 1 and 5 are optionally substituted. Furthermore, the pyridine compound employed in the method of the invention (hereafter Formula 3) is optionally substituted. The term "optionally substituted" in connection with these groups refers to groups that are unsubstituted or have at least one non-hydrogen substituent. Illustrative optional substituents include alkyl, alkenyl, cycloalkyl, cycloalkenyl, aryl, hydroxycarbonyl, formyl, alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, alkoxycarbonyl, hydroxy, alkoxy, alkenyloxy, alkynyloxy, cycloalkoxy, aryloxy, alkylthio, alkenylthio, alkynylthio, cycloalkylthio, arylthio, alkylsulfinyl, alkenylsulfinyl, alkynylsulfinyl, cycloalkylsulfinyl, arylsulfinyl, alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl, cycloalkylsulfonyl, arylsulfonyl, amino, alkylamino, alkenylamino, alkynylamino, arylamino, aminocarbonyl, alkylaminocarbonyl, alkenylaminocarbonyl, alkynylaminocarbonyl, arylaminocarbonyl, alkylaminocarbonyl, alkenylaminocarbonyl, alkynylaminocarbonyl, arylaminocarbonyloxy, alkoxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino and aryloxy-carbonylamino, silyl moieties and siloxy moieties, each further optionally substituted; halogen; cyano; and nitro. The optional further substituents are independently selected from groups like those illustrated above for the substituents themselves to give additional substituent groups for J and K such as haloalkyl, haloalkenyl and haloalkoxy. As another example, alkylamino can be further substituted with alkyl, giving dialkylamino. The substituents can also be tied together by figuratively removing one or two hydrogen atoms from each of two substituents or a substituent and the supporting molecular structure and joining the radicals to produce cyclic and polycyclic structures fused or appended to the molecular structure supporting the substituents. For example, tying together adjacent hydroxy and methoxy groups attached to, for example, a phenyl ring gives a fused dioxolane structure containing the linking group $-O-CH_2-O-$. Tying together a hydroxy group and the molecular structure to which it is attached can give cyclic ethers, including epoxides. Illustrative substituents also include oxygen, which when attached to carbon forms a carbonyl function or when attached to nitrogen forms an *N*-oxide.

One skilled in the art will appreciate that not all nitrogen-containing heterocycles can form *N*-oxides since the nitrogen requires an available lone pair for oxidation to the oxide; one skilled in the art will recognize those nitrogen containing heterocycles which can form *N*-oxides. One skilled in the art will also recognize that tertiary amines can form *N*-oxides. Synthetic methods for the preparation of *N*-oxides of heterocycles and tertiary amines are very well known by one skilled in the art including the oxidation of heterocycles and tertiary amines with peroxy acids such as peracetic and *m*-chloroperbenzoic acid (MCPBA), hydrogen peroxide, alkyl hydroperoxides such as *t*-butyl hydroperoxide, sodium perborate, and dioxiranes such as dimethyldioxirane. These methods for the preparation of *N*-oxides have been extensively described and reviewed in the literature, see for example:

T. L. Gilchrist in *Comprehensive Organic Synthesis*, vol. 7, pp 748–750, S. V. Ley, Ed., Pergamon Press; M. Tisler and B. Stanovnik in *Comprehensive Heterocyclic Chemistry*, vol. 3, pp 18–20, A. J. Boulton and A. McKillop, Eds., Pergamon Press; M. R. Grimmett and B. R. T. Keene in *Advances in Heterocyclic Chemistry*, vol. 43, pp 149–161, A. R. Katritzky, Ed., Academic Press; M. Tisler and B. Stanovnik in *Advances in Heterocyclic Chemistry*, vol. 9, pp 285–291, A. R. Katritzky and A. J. Boulton, Eds., Academic Press; and G. W. H. Cheeseman and E. S. G. Werstiuk in *Advances in Heterocyclic Chemistry*, vol. 22, pp 390–392, A. R. Katritzky and A. J. Boulton, Eds., Academic Press.

As referred to herein, "alkyl", used either alone or in compound words such as "alkylthio" or "haloalkyl" includes straight-chain or branched alkyl, such as, methyl, ethyl, *n*-propyl, *i*-propyl, and the different butyl, pentyl or hexyl isomers. "Alkenyl" includes straight-chain or branched alkenes such as ethenyl, 1-propenyl, 2-propenyl, and the different butenyl, pentenyl and hexenyl isomers. "Alkenyl" also includes polyenes such as 1,2-propadienyl and 2,4-hexadienyl. "Alkynyl" includes straight-chain or branched alkynes such as ethynyl, 1-propynyl, 2-propynyl and the different butynyl, pentynyl and hexynyl isomers. "Alkynyl" can also include moieties comprised of multiple triple bonds such as 2,5-hexadiynyl. "Alkoxy" includes, for example, methoxy, ethoxy, *n*-propyloxy, isopropyloxy and the different butoxy, pentoxy and hexyloxy isomers. "Alkenyloxy" includes straight-chain or branched alkenyloxy moieties. Examples of "alkenyloxy" include $\text{H}_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{O}$ and $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{O}$. "Alkynyloxy" includes straight-chain or branched alkynyloxy moieties. Examples of "alkynyloxy" include $\text{HC}\equiv\text{CCH}_2\text{O}$, $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{O}$ and $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}_2\text{O}$. "Alkylthio" includes branched or straight-chain alkylthio moieties such as methylthio, ethylthio, and the different propylthio, butylthio, pentylthio and hexylthio isomers. "Alkylsulfinyl" includes both enantiomers of an alkylsulfinyl group. Examples of "alkylsulfinyl" include $\text{CH}_3\text{S}(\text{O})$, $\text{CH}_3\text{CH}_2\text{S}(\text{O})$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{S}(\text{O})$, $(\text{CH}_3)_2\text{CHS}(\text{O})$ and the different butylsulfinyl, pentylsulfinyl and hexylsulfinyl isomers. Examples of "alkylsulfonyl" include $\text{CH}_3\text{S}(\text{O})_2$, $\text{CH}_3\text{CH}_2\text{S}(\text{O})_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{S}(\text{O})_2$, $(\text{CH}_3)_2\text{CHS}(\text{O})_2$ and the different butylsulfonyl, pentylsulfonyl and hexylsulfonyl isomers. "Alkylamino", "alkenylthio", "alkenylsulfinyl", "alkenylsulfonyl", "alkynylthio", "alkynylsulfinyl", "alkynylsulfonyl", and the like, are defined analogously to the above examples. Examples of "alkylcarbonyl" include $\text{C}(\text{O})\text{CH}_3$, $\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{CH}_3$ and $\text{C}(\text{O})\text{CH}(\text{CH}_3)_2$. Examples of "alkoxycarbonyl" include $\text{CH}_3\text{OC}(=\text{O})$, $\text{CH}_3\text{CH}_2\text{OC}(=\text{O})$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OC}(=\text{O})$, $(\text{CH}_3)_2\text{CHOC}(=\text{O})$ and the different butoxy- or pentoxycarbonyl isomers. "Cycloalkyl" includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. The term "cycloalkoxy" includes the same groups linked through an oxygen atom such as cyclopentyloxy and cyclohexyloxy. "Cycloalkylamino" means the amino nitrogen atom is attached to a cycloalkyl radical and a

hydrogen atom and includes groups such as cyclopropylamino, cyclobutylamino, cyclopentylamino and cyclohexylamino. "(Alkyl)(cycloalkyl)amino" means a cycloalkylamino group where the hydrogen atom is replaced by an alkyl radical; examples include groups such as (methyl)(cyclopropyl)amino, (butyl)(cyclobutyl)amino, (propyl)cyclopentylamino, (methyl)cyclohexylamino and the like. "Cycloalkenyl" includes groups such as cyclopentenyl and cyclohexenyl as well as groups with more than one double bond such as 1,3- and 1,4-cyclohexadienyl.

The term "halogen", either alone or in compound words such as "haloalkyl", includes fluorine, chlorine, bromine or iodine. Further, when used in compound words such as "haloalkyl", said alkyl may be partially or fully substituted with halogen atoms which may be the same or different. Examples of "haloalkyl" include F_3C , $ClCH_2$, CF_3CH_2 and CF_3CCl_2 .

The term "silyl moieties" refers to moieties containing at least one silicon atom bonded to the remainder of Formula 1 through said silicon atom and includes groups such as trialkylsilyl (examples include trimethylsilyl, triisopropylsilyl and dimethyl-*t*-butylsilyl) and dialkylarylsilyl groups (e.g. dimethylphenylsilyl). The term "siloxo moieties" refers to moieties containing at least one silicon atom bonded to an oxygen atom and connected to the remainder of Formula 1 through said oxygen atom and includes groups such as trialkylsiloxo (examples include trimethylsiloxo, triisopropylsiloxo and dimethyl-*t*-butylsiloxo) and dialkylarylsilyl groups (e.g. dimethylphenylsiloxo).

The total number of carbon atoms in a substituent group is indicated by the " C_i-C_j " prefix where *i* and *j* are, for example, numbers from 1 to 3; e.g., C_1-C_3 alkyl designates methyl through propyl.


There is no definite limit to the nature and size of the substituents on the pyridine of Formula 3, but the substituents are commonly alkyl, more particularly C_1-C_6 , more commonly C_1-C_4 , and most commonly C_1 (i.e. methyl). Typical examples of compounds of Formula 3 are pyridine, the picolines (i.e. 2-methylpyridine, 3-methylpyridine, 4-methylpyridine), the lutidines (e.g., 2,3-dimethylpyridine, 2,4-dimethylpyridine, 2,5-dimethylpyridine, 2,5-dimethylpyridine, 3,4-dimethylpyridine, 3,5-dimethylpyridine) and collidine. Other common substituents are dimethylamino (e.g., 4-(dimethylamino)pyridine) and pyrrolidino (e.g., 4-(pyrrolidino)pyridine). Furthermore, two substituents on Formula 3 can be tied together as already described to form other common pyridine derivatives, such as quinoline and isoquinoline. These quinoline and isoquinoline pyridine derivatives can also be further substituted.
































Although there is no definite limit to the size of J, optionally substituted alkyl moieties in J commonly include 1 to 6 carbon atoms, more commonly 1 to 4 carbon atoms, and most commonly 1 or 2 carbon atoms in the alkyl chain. Similarly, optionally substituted alkenyl

and alkynyl moieties in J commonly include 2 to 6 carbon atoms, more commonly 2 to 4 carbon atoms, and most commonly 2 or 3 carbon atoms in the alkenyl or alkynyl chain.


As indicated above, the carbon moiety J may be (among others) an aromatic ring or ring system. Examples of aromatic rings or ring systems include a phenyl ring, 5- or 6-membered heteroaromatic rings, aromatic 8-, 9- or 10-membered fused carbobicyclic ring systems and aromatic 8-, 9- or 10-membered fused heterobicyclic ring systems, wherein each ring or ring system is optionally substituted. The term "optionally substituted" in connection with these J groups refers to groups that are unsubstituted or have at least one non-hydrogen substituent. These carbon moieties may be substituted with as many optional substituents as can be accommodated by replacing a hydrogen atom with a non-hydrogen substituent on any available carbon or nitrogen atom. Commonly, the number of optional substituents (when present) ranges from one to four. An example of phenyl optionally substituted with from one to four substituents is the ring illustrated as U-1 in Exhibit 1, wherein R^V is any substituent and r is an integer from 0 to 4. Examples of aromatic 8-, 9- or 10-membered fused carbobicyclic ring systems optionally substituted with from one to four substituents include a naphthyl group optionally substituted with from one to four substituents illustrated as U-85 and a 1,2,3,4-tetrahydronaphthyl group optionally substituted with from one to four substituents illustrated as U-86 in Exhibit 1, wherein R^V is any substituent r is an integer from 0 to 4. Examples of 5- or 6-membered heteroaromatic rings optionally substituted with from one to four substituents include the rings U-2 through U-53 illustrated in Exhibit 1 wherein R^V is any substituent and r is an integer from 0 to 4. Examples of aromatic 8-, 9- or 10-membered fused heterobicyclic ring systems optionally substituted with from one to four substituents include U-54 through U-84 illustrated in Exhibit 1 wherein R^V is any substituent and r is an integer from 0 to 4. Other examples of J include a benzyl group optionally substituted with from one to four substituents illustrated as U-87 and a benzoyl group optionally substituted with from one to four substituents illustrated as U-88 in Exhibit 1, wherein R^V is any substituent and r is an integer from 0 to 4.

Although R^V groups are shown in the structures U-1 through U-85, it is noted that they do not need to be present since they are optional substituents. The nitrogen atoms that require substitution to fill their valence are substituted with H or R^V . Note that some U groups can only be substituted with less than 4 R^V groups (e.g. U-14, U-15, U-18 through U-21 and U-32 through U-34 can only be substituted with one R^V). Note that when the attachment point between $(R^V)_r$ and the U group is illustrated as floating, $(R^V)_r$ can be attached to any available carbon atom or nitrogen atom of the U group. Note that when the attachment point on the U group is illustrated as floating, the U group can be attached to the remainder of Formula I through any available carbon of the U group by replacement of a hydrogen atom.

Exhibit 1



12




As indicated above, the carbon moiety J may be (among others) saturated or partially saturated carbocyclic and heterocyclic rings, which can be further optionally substituted. The term "optionally substituted" in connection with these J groups refers to groups that are unsubstituted or have at least one non-hydrogen substituent. These carbon moieties may be substituted with as many optional substituents as can be accommodated by replacing a hydrogen atom with a non-hydrogen substituent on any available carbon or nitrogen atom.


Commonly, the number of optional substituents (when present) ranges from one to four. Examples of saturated or partially saturated carbocyclic rings include optionally substituted C₃-C₈ cycloalkyl and optionally substituted C₃-C₈ cycloalkenyl. Examples of saturated or partially saturated heterocyclic rings include optionally substituted 5- or 6-membered nonaromatic heterocyclic rings optionally including one or two ring members selected from the group consisting of C(=O), SO or S(O)₂. Examples of such J groups include those

illustrated as G-1 through G-35 in Exhibit 2. Note that when the attachment point on these G groups is illustrated as floating, the G group can be attached to the remainder of Formula 1 through any available carbon or nitrogen of the G group by replacement of a hydrogen atom. The optional substituents can be attached to any available carbon or nitrogen by replacing a hydrogen atom (said substituents are not illustrated in Exhibit 2 since they are optional substituents). Note that when G comprises a ring selected from G-24 through G-31, G-34 and G-35, Q² may be selected from O, S, NH or substituted N.


Exhibit 2



14



It is noted that the J group may be optionally substituted. As noted above, a J group may commonly comprise a U group or a G group further substituted with from one to four substituents. Thus J groups may comprise a U group or a G group selected from U-1 through U-88 or G-1 through G-35, and further substituted with additional substituents including from one to four U or G groups (which may be the same or different) and both the core U or G group and substituent U or G groups optionally further substituted. Of note are J groups comprising a U or G group substituted with one U or G group and optionally substituted with from one to three additional substituents. For example, J can be U-11, in which an R^V attached to the 1-nitrogen is the group U-41.



10

As noted above, K is, together with the two contiguous linking carbon atoms, a fused phenyl ring or a fused 5- or 6-membered heteroaromatic ring, each ring optionally substituted. The term "optionally substituted" in connection with these K rings refers to K rings that are unsubstituted or have at least one non-hydrogen substituent. An example of a K ring wherein the K ring is optionally substituted with from one to four R^t includes the ring system illustrated as K-38 (fused phenyl ring) in Exhibit 3 wherein n is an integer from 0 to 4 and R^t is any substituent. Example of said K rings wherein the K ring is optionally substituted with from one to three R^t include the ring systems K-1 to K-37 (5- or 6-membered heteroaromatic rings) in Exhibit 3, wherein n is an integer from 0 to 3 and R^t is any substituent. As with the carbon atoms in the ring, the nitrogen atoms that require substitution to fill their valence are substituted with hydrogen or with R^t . Although $(R^t)_n$ groups are shown in the structures K-1 to K-38, it is noted that R^t does not need to be present since it is an optional substituent. Note that some K rings can only be substituted with less than 3 R^t groups (e.g. K-7 through K-10, K-15, K-16, K-20, K-21, K-23, K-24, K-26 and K-27 can only be substituted with one R^t). In the exemplified K groups, the upper right bond is attached through the available linking carbon atom to the nitrogen atom of the oxazinone portion of Formula 1 and the lower right bond is attached through the available

linking carbon atom to the carbonyl atom of the oxazinone portion of Formula 1. The wavy line indicates that the K ring is attached to the remainder of Formula 1 as illustrated below.






Exhibit 3





Of note are K rings including optionally substituted thiophene, isoxazole, isothiazole, pyrazole, pyridine and pyrimidine rings. Of particular note are K rings K-1, K-14, K-15, K-18, K-23, K-28, K-29, K-30, K-31 and K-33, especially K-28, K-31 and K-33. Also of particular note is K-38 (optionally substituted phenyl).

Examples of optional substituents that can be attached to the U, G or K groups illustrated above include substituents selected from W. The K rings may also be substituted with the previously described optionally substituted U or optionally substituted G groups.

Each W is independently C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, C₃-C₆ cycloalkylamino, (C₁-C₄ alkyl)C₃-C₆ cycloalkylamino and C₃-C₆ trialkylsilyl.

Other suitable substituents include

B(OR¹⁷)₂; NH₂; SH; thiocyanato; C₃-C₈ trialkylsilyloxy; C₁-C₄ alkyldisulfide; SF₅; R¹⁹C(=E)-; R¹⁹C(=E)M-; R¹⁹MC(=E)-; (R¹⁹)MC(=E)M-; -OP(=Q)(OR¹⁹)₂; -SO₂MR¹⁹; R¹⁹SO₂M-;

wherein

each E is independently O, S, NR¹⁵, NOR¹⁵, NN(R¹⁵)₂, N-S=O, N-CN or N-NO₂;

each M is independently O, NR¹⁸ or S;

Q is O or S;

each R¹⁵ and each R¹⁹ is independently H; C₁-C₆ alkyl optionally substituted with

one or more substituents selected from the group consisting of CN, NO₂,

hydroxy, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl,

C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkylthio, C₁-C₄ haloalkylsulfinyl, C₁-C₄

haloalkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, CO₂H, C₂-C₆

alkoxycarbonyl, C₂-C₆ alkylcarbonyl, C₃-C₆ trialkylsilyl, and a phenyl ring

optionally substituted with one to three substituents independently selected from

W; C₁-C₆ haloalkyl; C₃-C₆ cycloalkyl; or a phenyl ring optionally substituted

with from one to three substituents independently selected from W;

each R¹⁷ is independently H or C₁-C₄ alkyl; or

B(OR¹⁷)₂ can form a ring wherein the two oxygen atoms are linked by a chain of two

to three carbons optionally substituted with one or two substituents

independently selected from methyl or C₂-C₆ alkoxycarbonyl; and

each R¹⁸ is independently H, C₁-C₆ alkyl or C₁-C₆ haloalkyl.

Of note are methods for preparing compounds of Formula 1 wherein J is C₁-C₆ alkyl,

C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkenyl, each optionally

substituted; or J is a phenyl ring, a benzyl group, a benzoyl group, a 5- or 6-membered

heteroaromatic ring, an aromatic 8-, 9- or 10-membered fused carbobicyclic ring system, an

aromatic 8-, 9- or 10-membered fused heterobicyclic ring system or a 5- or 6-membered

nonaromatic heterocyclic ring optionally including one or two ring members selected from

the group consisting of C(=O), SO or S(O)₂, each optionally substituted. Of particular note

are such methods wherein

K is, together with the two contiguous linking carbon atoms, a fused phenyl ring

optionally substituted with from one to four substituents independently selected

from G, U, W or R¹³; or a fused 5- or 6-membered heteroaromatic ring

optionally substituted with from one to three substituents independently selected

from G, U, W or R¹³;

J is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₈

cycloalkenyl, each optionally substituted with one or more substituents selected

from the group consisting of R¹², halogen, CN, NO₂, hydroxy, C₁-C₄ alkoxy,

C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈

dialkylamino, C₃-C₆ cycloalkylamino, and (C₁-C₄ alkyl)C₃-C₆ cycloalkylamino;

or

J is a phenyl ring, a benzyl group, a benzoyl group, a 5- or 6-membered heteroaromatic

ring, an aromatic 8-, 9- or 10-membered fused carbobicyclic ring system, an


- aromatic 8-, 9- or 10-membered fused heterobicyclic ring system or a 5- or 6-membered nonaromatic heterocyclic ring optionally including one or two ring members selected from the group consisting of C(=O), SO or S(O)₂, each optionally substituted with from one to four substituents independently selected from G, U, W or R¹³;
- 5 each G is a 5- or 6-membered nonaromatic heterocyclic ring optionally including one or two ring members selected from the group consisting of C(=O), SO or S(O)₂, each optionally substituted with from one to four substituents independently selected from W;
- 10 each U is a phenyl ring, a benzyl group, a benzoyl group, a 5- or 6-membered heteroaromatic ring, an aromatic 8-, 9- or 10-membered fused carbobicyclic ring system, an aromatic 8-, 9- or 10-membered fused heterobicyclic ring system, each optionally substituted with from one to four substituents independently selected from W;
- 15 each W is C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, C₃-C₆ cycloalkylamino, (C₁-C₄ alkyl)C₃-C₆ cycloalkylamino and C₃-C₆ trialkylsilyl;
- 20 each R¹² is independently R¹⁹C(=E)-; R¹⁹C(=E)L-; R¹⁹LC(=E)-; (R¹⁹)LC(=E)L-; -O(Q=)P(OR¹⁹)₂; -SO₂LR¹⁸; or R¹⁹SO₂L-;
- each R¹³ is B(OR¹⁷)₂; NH₂; SH; thiocyanato; C₃-C₈ trialkylsilyloxy; C₁-C₄ alkyldisulfide; SF₅; R¹⁹C(=E)-; R¹⁹C(=E)M-; R¹⁹MC(=E)-; (R¹⁹)MC(=E)M-;
- 25 -OP(=Q)(OR¹⁹)₂; -SO₂MR¹⁹; R¹⁹SO₂M-;
- each E is independently O, S, NR¹⁵, NOR¹⁵, NN(R¹⁵)₂, N-S=O, N-CN or N-NO₂;
- each M is independently O, NR¹⁸ or S;
- Q is O or S;
- each R¹⁵ and each R¹⁹ is independently H; C₁-C₆ alkyl optionally substituted with
- 30 one or more substituents selected from the group consisting of CN, NO₂, hydroxy, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkylthio, C₁-C₄ haloalkylsulfinyl, C₁-C₄ haloalkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, CO₂H, C₂-C₆ alkoxy carbonyl, C₂-C₆ alkyl carbonyl, C₃-C₆ trialkylsilyl, and a phenyl ring
- 35 optionally substituted with one to three substituents independently selected from W; C₁-C₆ haloalkyl; C₃-C₆ cycloalkyl; or a phenyl ring optionally substituted with from one to three substituents independently selected from W;
- each R¹⁷ is independently H or C₁-C₄ alkyl; or

$B(OR^{17})_2$ can form a ring wherein the two oxygen atoms are linked by a chain of two to three carbons optionally substituted with one or two substituents independently selected from methyl or C_2-C_6 alkoxycarbonyl; and each R^{18} is independently H, C_1-C_6 alkyl or C_1-C_6 haloalkyl.

5 Preferrably, K is, together with the two contiguous linking carbon atoms, a fused phenyl ring optionally substituted with from one to four substituents independently selected from W or R^{13} .

The compounds of Formula 1 can be prepared by one or more of the following methods and variations as described in Schemes 1-27. The definitions of J, K, L, M, R^4 , 10 through R^9 , R^v , X, Y and n in the compounds of Formulae 2-76 below are as defined above. Compounds of Formulae 1a, 2a-p and 5a-c are various subsets of the compounds of Formula 1, 2 and 5, respectively. Of note are compounds wherein K is selected from the group consisting of optionally substituted thiophene, isoxazole, isothiazole, pyrazole, pyridine and pyrimidine rings. Also of note are compounds wherein K is K-1, K-14, K-15, 15 K-18, K-23, K-28, K-29, K-30, K-31 and K-33. Of particular note are compounds of Formulae wherein K is K-28, K-31 and K-33. Also of particular note are compounds wherein K is an optionally substituted fused phenyl ring (K-38).


As shown in Scheme 1, according to the method of this invention a fused oxazinone of Formula 1 is prepared via coupling of a carboxylic acid of Formula 2 with an *ortho*-amino 20 carboxylic acid of Formula 5.



The reaction sequence (i.e. Steps (1), (2) and (3)) is typically conducted sequentially in the same reaction vessel and without isolation or purification of the products until the 25 overall reaction sequence is complete. Normally, Steps (1), (2) and (3) are all conducted in a suitable solvent. Typically the same solvent is used throughout the reaction sequence. The solvent(s) should be inert under the conditions of the reaction steps. Suitable solvents include aromatic hydrocarbons such as chlorobenzene or toluene, and preferred solvents include esters such as ethyl acetate or butyl acetate; ketones such as acetone, 2-butanone, or 30 4-methyl-2-pentanone; ethers such as tetrahydrofuran (THF) or dioxane; nitriles such as

acetonitrile; and halocarbons such as dichloromethane or chloroform. More preferred solvents include acetonitrile, ethyl acetate, acetone, THF and dichloromethane. Most preferred are acetonitrile and acetone. The overall reaction sequence is typically conducted at temperatures ranging from -30°C to $+50^{\circ}\text{C}$. Preferred is the addition to the reaction vessel (charging) of all components at temperatures from -10°C to $+5^{\circ}\text{C}$, with reaction periods of from 1 to 60 minutes (preferred are reaction periods of from 5 to 15 minutes) between additions. Except where noted otherwise, the components can be added as mixtures, followed by warming to from $+20^{\circ}\text{C}$ to $+30^{\circ}\text{C}$ for 0.5 to 24 hours (preferably 2 to 4 hours). Charging of a component (for example, a reactant, a solvent, etc.) means adding the component during the step. One of ordinary skill in the art will recognize that a component may be added (i.e. charged) in various ways, for example as a batch, intermittent or continuous feed depending on process design. The manner in which a component is added and/or reacted can influence the actual mole ratio of that component to others over the duration of the step. However, the overall amount of a component added during a step is considered to be the amount charged, and is used herein for determining nominal mole ratios.

In Step (1) a carboxylic acid of Formula 2




is contacted with a sulfonyl chloride of Formula 4



4

in the presence of an optionally substituted pyridine compound of Formula 3. The nominal mole ratio (i.e. the ratio of reactants charged during Step (1)) of the Formula 4 compound to the Formula 2 compound is typically from about 1.0 to 1.5, and preferably is from about 1.1 to 1.3. Preferred Formula 4 compounds include methanesulfonyl chloride, propanesulfonyl chloride and benzenesulfonyl chloride. Methanesulfonyl chloride is more preferred for reasons of lower cost and/or less waste. The nominal mole ratio of the Formula 3 compound charged in Step (1) to the Formula 2 compound charged in Step (1) is typically from about 1.0 to 2.0, and is preferably from about 1.4 to 1.7. Preferred Formula 3 compounds include 2-picoline, 3-picoline, 2,6-lutidine, and pyridine. The compounds of Formulae 2, 3 and 4 may be combined in any order except that compound 2 cannot be added to the reaction mixture last. A preferred order of addition is to add a mixture of Formula 2 and Formula 3 in a solvent to a solution of Formula 4 in the same solvent.


In Step (2), the mixture prepared in (1) is contacted with an anthranilic acid of Formula 5



in the presence of an optionally substituted pyridine compound of Formula 3. Typically, the optionally substituted pyridine compound of Formula 3 in Step (2) is the same compound as that used in Step (1). The nominal mole ratio of the Formula 5 compound used in Step (2) to the Formula 2 compound charged in Step (1) is typically from about 0.9 to 1.1, and is preferably about 1.0. Normally, a second portion of Formula 3 compound is charged during Step (2); and the nominal mole ratio of the second portion of Formula 3 compound charged in Step (2) to the Formula 2 compound charged in Step (1) is typically from about 2.0 to 4.0, and preferably is from about 2.9 to 3.5. The compounds of Formulae 5 and 3 can be added to the mixture prepared in Step (1) in either order or as a mixture in solvent.

In step (3) the mixture prepared in Step (2) is contacted with a sulfonyl chloride of Formula 4. Typically, the sulfonyl chloride in Step (3) is the same compound as that used in Step (1). The nominal mole ratio of the second portion of the Formula 4 compound charged in Step (3) to the Formula 2 compound charged in Step (1) is typically from about 1.0 to 1.5, and preferably is from about 1.1 to 1.3. Typically, Formula 4 is added to the mixture in the reaction solvent.

Preferred methods of this invention include the method wherein the carboxylic acid of Formula 2 is Formula 2', the *ortho*-amino carboxylic acid of Formula 5 is Formula 5' and the product compound of Formula 1 is Formula 1a



20

wherein

X is N or CR^6 ;

Y is N or CH;

R⁴ is C₁-C₄ alkyl or halogen;

R⁵ is H, C₁-C₄ alkyl, C₁-C₄ haloalkyl or halogen;

R⁶ and R⁷ are independently H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, halogen, CN or C₁-C₄ haloalkoxy;


R⁸ is H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, C₃-C₆ cycloalkylamino, (C₁-C₄ alkyl)(C₃-C₆ cycloalkyl)amino, C₂-C₄ alkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₂-C₆ alkylaminocarbonyl, C₃-C₈ dialkylaminocarbonyl or C₃-C₆ trialkylsilyl; and

R⁹ is CF₃, OCF₃, OCHF₂, OCH₂CF₃, S(O)_pCF₃, S(O)_pCHF₂ or halogen.

Compounds of Formula 1a may be prepared in this way and used for preparing compounds of Formula III.


Ortho-amino carboxylic acid esters of Formula 5 (Formula 7) can be prepared from monoesters of *ortho*-dicarboxylic acids of Formula 6 via rearrangement of the corresponding acyl azide and hydrolysis of the resulting isocyanate (or alternatively by trapping of the isocyanate with an alcohol and cleaving of the resulting carbamate) as shown in Scheme 2.

Scheme 2



Alternatively *ortho*-amino carboxylic acid esters of Formula 7 can be prepared from *ortho*-carboxamide carboxylic esters of Formula 8 by Hoffman rearrangement with reagents such as sodium hydroxide and bromine as shown in Scheme 3.


Scheme 3




Compounds of Formulae 6 and 8 are known in the art or can be readily prepared from compounds known in the art. (For example, see *Tetrahedron*, 1997, 53, 14497; *J. Chem. Soc., Perkin Trans. 1*, 1996, 10, 1035; PCT Patent Publication WO 92/08724 and European Patent Application EP 418667).

5 Carboxylic ester compounds can be converted to carboxylic acid compounds by numerous methods including nucleophilic cleavage under anhydrous conditions or hydrolytic methods involving the use of either acids or bases (see T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 2nd ed., John Wiley & Sons, Inc., New York, 1991, pp. 224-269 for a review of methods). In many cases, base-catalyzed hydrolytic
 10 methods are preferred. Suitable bases include alkali metal (such as lithium, sodium or potassium) hydroxides. For example, the ester can be dissolved in a mixture of water and an alcohol such as ethanol. Upon treatment with sodium hydroxide or potassium hydroxide, the ester is saponified to provide the sodium or potassium salt of the carboxylic acid. Acidification with a strong acid, such as hydrochloric acid or sulfuric acid, yields the
 15 carboxylic acid of Formula 2f. The carboxylic acid can be isolated by methods known to those skilled in the art, including crystallization, extraction and distillation.

Anthranilic acids (or esters thereof) of Formula 5a (compounds of Formula 5 wherein K is a fused phenyl ring) are typically available from the corresponding 2-nitrobenzoic acids (or esters) of Formula 9 via catalytic hydrogenation of the nitro group (Scheme 4). Typical
 20 procedures involve reduction with hydrogen in the presence of a metal catalyst such as palladium on carbon or platinum oxide in hydroxylic solvents such as ethanol and isopropanol. They can also be prepared by reduction with zinc in acetic acid. These methods for reducing nitro groups are well documented in the chemical literature.

Scheme 4

As shown in Scheme 5, anthranilic acids containing an R^5 substituent of chloro, bromo or iodo can be prepared by direct halogenation of an anthranilic acid of Formula **5b** with *N*-chlorosuccinimide (NCS), *N*-bromosuccinimide (NBS) or *N*-iodosuccinimide (NIS) respectively in solvents such as *N,N*-dimethylformamide (DMF) to produce the corresponding substituted acid of Formula **5c**.


Scheme 5

Benzoic acids of Formula 2 (*J* is optionally substituted phenyl) are generally well known in the art as are procedures for their preparation.

Benzoic acids of Formula 2a may be prepared from the benzonitriles of Formula 10 by hydrolysis (Scheme 6). The conditions used may involve the use of a base such as an alkaline metal hydroxide or alkoxide (e.g. potassium or sodium hydroxide) in a solvent such as water, ethanol or ethylene glycol (see e.g. *J. Chem. Soc.* 1948, 1025). Alternatively, the hydrolysis may be carried out using an acid such as sulfuric acid or phosphoric acid in a suitable solvent such as water (see e.g. *Org. Synth.* 1955, Coll vol. 3, 557). The choice of the conditions is contingent on the stability of any optional substituents present on the aromatic ring to the reaction conditions and elevated temperatures are usually employed to achieve this transformation.

25


Scheme 6



Nitriles of Formula 10 may be prepared from anilines of Formula 11 by the classical sequence involving diazotization and treatment of the intermediate diazonium salt with a copper cyanide salt (see e.g. *J. Amer. Chem. Soc.* 1902, 24, 1035).

5

Scheme 7




Certain heterocyclic acids of Formula 2, wherein J is an optionally substituted heterocycle, can be prepared by procedures outlined in Schemes 8 through 27. Both general and specific references to a wide variety of heterocyclic acids including thiophenes, furans, pyridines, pyrimidines, triazoles, imidazoles, pyrazoles, thiazoles, oxazoles, isothiazoles, thiadiazoles, oxadiazoles, triazines, pyrazines, pyridazines, and isoxazoles can be found in the following compendia: *Rodd's Chemistry of Carbon Compounds*, Vol. IVa to IVI., S. Coffey editor, Elsevier Scientific Publishing, New York, 1973; *Comprehensive Heterocyclic Chemistry*, Vol. 1-7, A. R. Katritzky and C. W. Rees editors, Pergamon Press, New York, 1984; *Comprehensive Heterocyclic Chemistry II*, Vol. 1-9, A. R. Katritzky, C. W. Rees, and E. F. Scriven editors, Pergamon Press, New York, 1996; and the series, *The Chemistry of Heterocyclic Compounds*, E. C. Taylor, editor, Wiley, New York. Noteworthy heterocyclic acids suitable for use in this invention include pyridine acids, pyrimidine acids, pyrazole acids and pyrrole acids. Procedures for the synthesis of representative examples of each are detailed in Schemes 8 through 27. A variety of heterocyclic acids and general methods for their synthesis may be found in PCT Patent Publication WO 98/57397.

The synthesis of representative pyridine acids of Formula 2b is depicted in Scheme 8. This procedure involves the known synthesis of pyridines from β -ketoesters (Formula 16) and 4-aminobutenones (Formula 15). Substituent groups R^a and R^b include e.g. alkyl, haloalkyl, and optionally substituted aromatic and heteroaromatic rings.


26

Scheme 8



The synthesis of representative pyrimidine acids (Formula 2c) is depicted in Scheme 9. This procedure involves the known synthesis of pyrimidines from vinylidene- β -ketoesters (Formula 16) and amidines (Formula 20). Substituent groups R^a and R^b include e.g. alkyl, haloalkyl, and optionally substituted aromatic and heteroaromatic rings.


Scheme 9



Syntheses of representative pyrazole acids (Formula 2d-2g) are depicted in Schemes 10 through 13. The synthesis of 2d in Scheme 10 involves as the key step introduction of the R^a substituent via alkylation of the pyrazole. The alkylating agent $R^a\text{-Lg}$ (wherein Lg is a leaving group such as Cl, Br, I, sulfonates such as p-toluenesulfonate or methanesulfonate or sulfates such as $-\text{SO}_2\text{OR}^5(\text{c})$) includes R^a groups such as $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_6$ cycloalkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_2\text{-C}_6$ haloalkenyl, $\text{C}_2\text{-C}_6$ haloalkynyl,


C₃-C₆ halocycloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₃-C₈ dialkylamino-carbonyl, C₃-C₆ trialkylsilyl; or phenyl, benzyl, benzoyl, 5- or 6-membered heteroaromatic rings or aromatic 8-, 9- or 10-membered fused heterobicyclic ring systems, each ring or ring system optionally substituted. Oxidation of the methyl group affords the pyrazole carboxylic acid. Some notable R^b groups include haloalkyl.

Scheme 10



Some pyrazole acids of Formula 2d may be prepared via metallation and carboxylation of pyrazoles of Formula 26 as the key step (Scheme 11). This reaction is typically conducted by treating compounds of Formula 25 with lithium diisopropylamide (LDA) to form an anion and then contacting the anion with carbon dioxide. The R^a group is introduced in a manner similar to that of Scheme 13, i.e. via alkylation with an R^a alkylating agent. Representative R^b groups include e.g. cyano and haloalkyl.


Scheme 11



This procedure is particularly useful for preparing 1-(2-pyridinyl)pyrazolecarboxylic acids of Formula 2e, wherein R^a is a substituted 2-pyridinyl ring, as shown in Scheme 12. Reaction of a pyrazole of Formula 27 with a 2-halopyridine of Formula 28 affords good yields of the 1-pyridinylpyrazole of Formula 29 with good specificity for the desired regiochemistry. Metallation of 29 with LDA followed by quenching of the lithium salt with carbon dioxide affords the 1-(2-pyridinyl)pyrazolecarboxylic acid of Formula 2e.

28

Scheme 15




wherein R^7 is H, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen, CN or C_1 - C_4 haloalkoxy; and

R^9 is CF_3 , OCF_3 , OCHF_2 , OCH_2CF_3 , $\text{S(O)}_p\text{CF}_3$, $\text{S(O)}_p\text{CHF}_2$ or halogen

5 Other pyrazoles of Formula 2d can be prepared via reaction of hydrazine of Formula 31 with a pyruvate of Formula 30 to yield pyrazole esters of Formula 32 (Scheme 13). Hydrolysis of the ester affords the pyrazole acids 2d. This procedure is particularly useful for the preparation of compounds wherein R^a is optionally substituted phenyl and R^b is haloalkyl.

10

Scheme 13




15 Pyrazole acids of Formula 2d can also be prepared via 3+2 cycloaddition of an appropriately substituted nitrilimine of Formula 33 with either substituted propiolates of Formula 34 or acrylates of Formula 35 (Scheme 14). Cycloaddition with acrylates requires additional oxidation of the intermediate pyrazoline to the pyrazole. Hydrolysis of the ester of Formula 36 affords the pyrazole acids 2d. Preferred iminohalides for this reaction include the trifluoromethyl iminochloride (37) and the iminodibromide (38). Compounds such as 37 are known (*J. Heterocycl. Chem.* 1985, 22(2), 565-8). Compounds such as 38 are available by known methods (*Tetrahedron Letters* 1999, 40, 2605). These procedures are particularly useful for the preparation of compounds where R^a is optionally substituted phenyl and R^b is haloalkyl or bromo.

20

29


Scheme 14




The starting pyrazoles of Formula 25 are known compounds or can be prepared according to known methods. The pyrazole of Formula 25a (the compound of Formula 25 wherein R^b is CF_3) can be prepared by literature procedures (*J. Fluorine Chem.* 1991, 53(1), 61-70). The pyrazoles of Formula 25b (compounds of Formula 25 wherein R^b is Cl or Br) can be prepared by literature procedures (*Chem. Ber.* 1966, 99(10), 3350-7). A useful alternative method for the preparation of a compound of Formula 25b is depicted in Scheme 15. Metallation of the sulfamoyl pyrazole of Formula 39 with *n*-butyllithium followed by direct halogenation of the anion with either hexachloroethane (for R^b being Cl) or 1,2-dibromotetrachloroethane (for R^b being Br) affords the halogenated derivatives of Formula 40. Removal of the sulfamoyl group with trifluoroacetic acid (TFA) at room temperature proceeds cleanly and in good yield to afford the pyrazoles of Formula 25c. One skilled in the art will recognize that Formula 25c is a tautomer of Formula 25b.

15

Scheme 15



Pyrazolecarboxylic acids of Formula 2f wherein R¹⁰ is CF₃ can be prepared by the method outlined in Scheme 16.



- 5 Reaction of a compound of Formula 41 wherein R¹⁰ is C₁-C₄ alkyl with a suitable base in a suitable organic solvent affords the cyclized product of Formula 42 after neutralization with an acid such as acetic acid. The suitable base can be, for example but not limitation, sodium hydride, potassium *t*-butoxide, dimethyl sodium (CH₃S(O)CH₂⁻ Na⁺), alkali metal (such as lithium, sodium or potassium) carbonates or hydroxides, tetraalkyl (such as methyl, ethyl or
- 10 butyl)ammonium fluorides or hydroxides, or 2-*tert*-butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphonine. The suitable organic solvent can be, for example but not limitation, acetone, acetonitrile, tetrahydrofuran, dichloromethane, dimethylsulfoxide, or *N,N*-dimethylformamide. The cyclization reaction is usually conducted in a temperature range from about 0 to 120 °C. The effects of solvent, base, temperature and addition time
- 15 are all interdependent, and choice of reaction conditions is important to minimize the formation of byproducts. A preferred base is tetrabutyl-ammonium fluoride.

Dehydration of the compound of Formula 42 to give the compound of Formula 43, followed by converting the carboxylic ester function to carboxylic acid, affords the compound of Formula 2f. The dehydration is effected by treatment with a catalytic amount


20 of a suitable acid. This catalytic acid can be, for example but not limitation, sulfuric acid. The reaction is generally conducted using an organic solvent. As one skilled in the art will realize, dehydration reactions may be conducted in a wide variety of solvents in a temperature range generally between about 0 and 200 °C, more preferably between about 0 and 100 °C). For the dehydration in the method of Scheme 16, a solvent comprising acetic acid and temperatures of about 65 °C are preferred. Carboxylic ester compounds can be

25 converted to carboxylic acid compounds by numerous methods including nucleophilic cleavage under anhydrous conditions or hydrolytic methods involving the use of either acids or bases (see T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 2nd ed., John Wiley & Sons, Inc., New York, 1991, pp. 224-269 for a review of methods). For

the method of Scheme 16, base-catalyzed hydrolytic methods are preferred. Suitable bases include alkali metal (such as lithium, sodium or potassium) hydroxides. For example, the ester can be dissolved in a mixture of water and an alcohol such as ethanol. Upon treatment with sodium hydroxide or potassium hydroxide, the ester is saponified to provide the sodium or potassium salt of the carboxylic acid. Acidification with a strong acid, such as hydrochloric acid or sulfuric acid, yields the carboxylic acid of Formula 2f. The carboxylic acid can be isolated by methods known to those skilled in the art, including crystallization, extraction and distillation.

Compounds of Formula 41 can be prepared by the method outlined in Scheme 17.

10




wherein R^9 is CF_3 and R^{10} is C_1-C_4 alkyl.

Treatment of a hydrazine compound of Formula 44 with a ketone of Formula 45 in a solvent such as water, methanol or acetic acid gives the hydrazone of Formula 46. One skilled in the art will recognize that this reaction may require catalysis by an optional acid and may also require elevated temperatures depending on the molecular substitution pattern of the hydrazone of Formula 46. Reaction of the hydrazone of Formula 46 with the compound of Formula 47 in a suitable organic solvent such as, for example but not limitation, dichloromethane or tetrahydrofuran in the presence of an acid scavenger such as triethylamine provides the compound of Formula 41. The reaction is usually conducted at a temperature between about 0 and 100 °C. Hydrazine compounds of Formula 44 can be prepared by standard methods, such as by contacting the corresponding halo compound of Formula 28 (Scheme 12) with hydrazine.

Pyrazolecarboxylic acids of Formula 2g wherein R^9 is Cl or Br can be prepared by the method outlined in Scheme 18.

32

Scheme 18




wherein R^{10} is C_1 - C_4 alkyl.

Oxidization of the compound of Formula 48 optionally in the presence of acid to give the compound of Formula 49 followed by conversion of the carboxylic ester function to the carboxylic acid provides the compound of Formula 2g. The oxidizing agent can be hydrogen peroxide, organic peroxides, potassium persulfate, sodium persulfate, ammonium persulfate, potassium monopersulfate (e.g., Oxone®) or potassium permanganate. To obtain complete conversion, at least one equivalent of oxidizing agent versus the compound of Formula 48 should be used, preferably between about one to two equivalents. This oxidation is typically carried out in the presence of a solvent. The solvent can be an ether, such as tetrahydrofuran, *p*-dioxane and the like, an organic ester, such as ethyl acetate, dimethyl carbonate and the like, or a polar aprotic organic such as *N,N*-dimethylformamide, acetonitrile and the like. Acids suitable for use in the oxidation step include inorganic acids, such as sulfuric acid, phosphoric acid and the like, and organic acids, such as acetic acid, benzoic acid and the like. The acid, when used, should be used in greater than 0.1 equivalents versus the compound of Formula 48. To obtain complete conversion, one to five equivalents of acid can be used. The preferred oxidant is potassium persulfate and the oxidation is preferably carried out in the presence of sulfuric acid. The reaction can be carried out by mixing the compound of Formula 48 in the desired solvent and, if used, the acid. The oxidant can then be added at a convenient rate. The reaction temperature is typically varied from as low as about 0 °C up to the boiling point of the solvent in order to obtain a reasonable reaction time to complete the reaction, preferably less than 8 hours. The desired product, a compound of Formula 49, can be isolated by methods known to those skilled in the art, including crystallization, extraction and distillation. Methods suitable for converting the ester of Formula 49 to the carboxylic acid of Formula 2g are already described for Scheme 16.

Compounds of Formula 48 can be prepared from corresponding compounds of Formula 50 as shown in Scheme 19.

Scheme 19



wherein R^{10} is C_1 - C_4 alkyl.

Treatment of a compound of Formula 50 with a halogenating reagent, usually in the presence of a solvent, affords the corresponding halo compound of Formula 48. Halogenating reagents that can be used include phosphorus oxyhalides, phosphorus trihalides, phosphorus pentahalides, thionyl chloride, dihalotrialkylphosphoranes, dihalodiphenylphosphoranes, oxalyl chloride and phosgene. Preferred are phosphorus oxyhalides and phosphorus pentahalides. To obtain complete conversion, at least 0.33 equivalents of phosphorus oxyhalide versus the compound of Formula 50 should be used (i.e. the mole ratio of phosphorus oxyhalide to the compound of Formula 50 is 0.33), preferably between about 0.33 and 1.2 equivalents. To obtain complete conversion, at least 0.20 equivalents of phosphorus pentahalide versus the compound of Formula 50 should be used, preferably between about 0.20 and 1.0 equivalents. Compounds of Formula 50 wherein R^{10} is C_1 - C_4 alkyl are preferred for this reaction. Typical solvents for this halogenation include halogenated alkanes, such as dichloromethane, chloroform, chlorobutane and the like, aromatic solvents, such as benzene, xylene, chlorobenzene and the like, ethers, such as tetrahydrofuran, *p*-dioxane, diethyl ether, and the like, and polar aprotic solvents such as acetonitrile, *N,N*-dimethylformamide, and the like. Optionally, an organic base, such as triethylamine, pyridine, *N,N*-dimethylaniline or the like, can be added. Addition of a catalyst, such as *N,N*-dimethylformamide, is also an option. Preferred is the process in which the solvent is acetonitrile and a base is absent. Typically, neither a base nor a catalyst is required when acetonitrile solvent is used. The preferred process is conducted by mixing the compound of Formula 50 in acetonitrile. The halogenating reagent is then added over a convenient time, and the mixture is then held at the desired temperature until the reaction is complete. The reaction temperature is typically between 20 °C and the boiling point of acetonitrile, and the reaction time is typically less than 2 hours. The reaction mass is then neutralized with an inorganic base, such as sodium bicarbonate, sodium hydroxide and the like, or an organic base, such as sodium acetate. The desired product, a compound of

Formula 48, can be isolated by methods known to those skilled in the art, including crystallization, extraction and distillation.

Alternatively, compounds of Formula 48 wherein R^9 is Br or Cl can be prepared by treating the corresponding compounds of Formula 48 wherein R^9 is a different halogen (e.g., Cl for making Formula 48 wherein R^9 is Br) or a sulfonate group such as *p*-toluenesulfonate with hydrogen bromide or hydrogen chloride, respectively. By this method the R^9 halogen or sulfonate substituent on the Formula 48 starting compound is replaced with Br or Cl from hydrogen bromide or hydrogen chloride, respectively. The reaction is conducted in a suitable solvent such as dibromomethane, dichloromethane or acetonitrile. The reaction can be conducted at or near atmospheric pressure or above atmospheric pressure in a pressure vessel. When R^9 in the starting compound of Formula 48 is a halogen such as Cl, the reaction is preferably conducted in such a way that the hydrogen halide generated from the reaction is removed by sparging or other suitable means. The reaction can be conducted between about 0 and 100 °C, most conveniently near ambient temperature (e.g., about 10 to 40 °C), and more preferably between about 20 and 30 °C. Addition of a Lewis acid catalyst (such as aluminum tribromide for preparing Formula 48 wherein R^9 is Br) can facilitate the reaction. The product of Formula 48 is isolated by the usual methods known to those skilled in the art, including extraction, distillation and crystallization.


Starting compounds of Formula 48 wherein R^9 is Cl or Br can be prepared from corresponding compounds of Formula 50 as already described. Starting compounds of Formula 48 wherein R^9 is a sulfonate group can likewise be prepared from corresponding compounds of Formula 50 by standard methods such as treatment with a sulfonyl chloride (e.g., *p*-toluenesulfonyl chloride) and base such as a tertiary amine (e.g., triethylamine) in a suitable solvent such as dichloromethane.

Pyrazolecarboxylic acids of Formula 2h wherein R^9 is OCH_2CF_3 or Formula 2i wherein R^9 is $OCHF_2$ can be prepared by the method outlined in Scheme 20. In this method, instead of being halogenated as shown in Scheme 19, the compound of Formula 50 is oxidized to the compound of Formula 51. The reaction conditions for this oxidation are as already described for the conversion of the compound of Formula 48 to the compound of Formula 49 in Scheme 18.

The compound of Formula 51 is then alkylated to form the compound of Formula 54 (R^9 is OCH_2CF_3) by contact with an alkylating agent CF_3CH_2Lg (52) in the presence of a base. In the alkylating agent 52, Lg is a nucleophilic reaction leaving group such as halogen (e.g., Br, I), $OS(O)_2CH_3$ (methanesulfonate), $OS(O)_2CF_3$, $OS(O)_2Ph$ -*p*-CH₃ (*p*-toluenesulfonate), and the like; methanesulfonate works well. The reaction is conducted in the presence of at least one equivalent of a base. Suitable bases include inorganic bases, such as alkali metal (such as lithium, sodium or potassium) carbonates and hydroxides, and organic bases, such as triethylamine, diisopropylethylamine and 1,8-diazabicyclo-

[5.4.0]undec-7-ene. The reaction is generally conducted in a solvent, which can comprise alcohols, such as methanol and ethanol, halogenated alkanes, such as dichloromethane, aromatic solvents, such as benzene, toluene and chlorobenzene, ethers, such as tetrahydrofuran, and polar aprotic solvents, such as acetonitrile, *N,N*-dimethylformamide, and the like. Alcohols and polar aprotic solvents are preferred for use with inorganic bases. Potassium carbonate as base and acetonitrile as solvent are preferred. The reaction is typically conducted between about 0 and 150 °C, and more typically between ambient temperature and 100 °C.

Scheme 20




wherein R²⁰ is C₁-C₄ alkyl, and Lg is a leaving group.

The compound of Formula 51 can also be alkylated to form the compound of Formula 55 (R⁹ is OCHF₂) by contact with difluorocarbene, prepared from CHClF₂ (53) in the presence of a base. The reaction is generally conducted in a solvent, which can comprise ethers, such as tetrahydrofuran or dioxane, and polar aprotic solvents, such as acetonitrile, *N,N*-dimethylformamide, and the like. The base can be selected from inorganic bases such as potassium carbonate, sodium hydroxide or sodium hydride. Preferably the reaction is conducted using potassium carbonate with *N,N*-dimethylformamide as the solvent. The product of Formula 54 or 55 can be isolated by conventional techniques such as extraction. The esters can then be converted to the carboxylic acids of Formula 2h or 2i by the methods already described for the conversion of Formula 43 to Formula 2f in Scheme 16.

Compounds of Formula 50 can be prepared from compounds of Formula 44 (see Scheme 17) as outlined in Scheme 21.

Scheme 21



5 wherein R¹⁰ is C₁-C₄ alkyl.


In this method, a hydrazine compound of Formula 44 is contacted with a compound of Formula 56 (a fumarate ester or maleate ester or a mixture thereof may be used) in the presence of a base and a solvent. The base is typically a metal alkoxide salt, such as sodium methoxide, potassium methoxide, sodium ethoxide, potassium ethoxide, potassium *tert*-butoxide, lithium *tert*-butoxide, and the like. Greater than 0.5 equivalents of base versus the compound of Formula 56 should be used, preferably between 0.9 and 1.3 equivalents. Greater than 1.0 equivalents of the compound of Formula 44 should be used, preferably between 1.0 to 1.3 equivalents. Polar protic and polar aprotic organic solvents can be used, such as alcohols, acetonitrile, tetrahydrofuran, *N,N*-dimethylformamide, dimethyl sulfoxide and the like. Preferred solvents are alcohols such as methanol and ethanol. It is especially preferred that the alcohol be the same as that making up the fumarate or maleate ester and the alkoxide base. The reaction is typically conducted by mixing the compound of Formula 56 and the base in the solvent. The mixture can be heated or cooled to a desired temperature and the compound of Formula 44 added over a period of time. Typically reaction temperatures are between 0 °C and the boiling point of the solvent used. The reaction may be conducted under greater than atmospheric pressure in order to increase the boiling point of the solvent. Temperatures between about 30 and 90 °C are typically preferred. The addition time can be as quick as heat transfer allows. Typical addition times are between 1 minute and 2 hours. Optimum reaction temperature and addition time vary depending upon the identities of the compounds of Formula 44 and Formula 56. After addition, the reaction mixture can be held for a time at the reaction temperature. Depending upon the reaction temperature, the required hold time may be from 0 to 2 hours. Typical hold times are 10 to 60 minutes. The reaction mass then can be acidified by adding an organic acid, such as acetic acid and the like, or an inorganic acid, such as hydrochloric acid, sulfuric acid and the like. Depending on the reaction conditions and the means of isolation, the -CO₂R¹⁰ function on the compound of Formula 50 may be hydrolyzed to -CO₂H; for example, the presence of water in the reaction mixture can promote such hydrolysis. If the

carboxylic acid ($-\text{CO}_2\text{H}$) is formed, it can be converted back to $-\text{CO}_2\text{R}^{10}$ wherein R^{10} is $\text{C}_1\text{-C}_4$ alkyl using esterification methods well known in the art. The desired product, a compound of Formula 50, can be isolated by methods known to those skilled in the art, such as crystallization, extraction or distillation.

- 5 The synthesis of representative pyrazole acids of Formula 2j is depicted in Scheme 22. Reaction of a dimethylaminoylidene ketoester of Formula 58 with substituted hydrazines of formula 31 affords the pyrazoles of Formula 59. Notable R^a substituents include alkyl and haloalkyl, with 2,2,2-trifluoroethyl especially noteworthy. The esters of Formula 59 are converted to the acids of Formula 2j by standard hydrolysis methods.

10


Scheme 22



The synthesis of pyrazole acids of Formula 2k, wherein R^b is a substituted 2-pyridyl moiety attached to the 5-position of the pyrazole ring, is depicted in Scheme 23. This synthesis is conducted according to the general synthesis described in Scheme 22.


15

Scheme 23



The synthesis of representative pyrazole acids of Formula 2m, as well as an alternative synthesis of Formula 2j, is depicted in Scheme 24. Reaction of the dimethylaminoylidene ketoester of Formula 58 with hydrazine affords the pyrazole of Formula 63. Reaction of the pyrazole 63 with alkylating agents of Formula 23 (R^a -Lg wherein Lg is a leaving group such as halogen (e.g., Br, I), $OS(O)_2CH_3$ (methanesulfonate), $OS(O)_2CF_3$, $OS(O)_2Ph$ -*p*-CH₃ (*p*-toluenesulfonate), and the like) affords a mixture of pyrazoles of Formulae 64 and 65. This mixture of pyrazole isomers is readily separated by chromatographic methods and converted to the corresponding acids 2m and 2j. Noteworthy R^a substituents include alkyl and haloalkyl groups.


Scheme 24



Of note is the synthesis of pyridinylpyrazole acids of Formula 2n, wherein R^b is a substituted 2-pyridinyl and attached to the 3-position of the pyrazole ring, as well as an alternative synthesis of Formula 2k, is depicted in Scheme 25. This synthesis is conducted according to the general synthesis described in Scheme 24.


39

Scheme 25




A general synthesis of pyrrole acids of Formula 2o is depicted in Scheme 26. Treatment of a compound of Formula 69 with 2,5-dimethoxytetrahydrofuran (70) affords a pyrrole of Formula 71. Formylation of the pyrrole 71 to provide the aldehyde of Formula 72 can be accomplished by using standard Vilsmeier-Haack formylation conditions, such as *N,N*-dimethylformamide (DMF) and phosphorus oxychloride. Halogenation of the compound of Formula 72 with *N*-halosuccinimides (NXS) such as *N*-chlorosuccinimide or *N*-bromosuccinimide occurs preferentially at the 4-position of the pyrrole ring. Oxidation of the halogenated aldehyde affords the pyrrole acid of Formula 2o. The oxidation can be accomplished by using a variety of standard oxidation conditions.

Scheme 26



The synthesis of certain pyridinylpyrrole acids of Formula 2p is depicted in Scheme 27. The compound of Formula 74, 3-chloro-2-aminopyridine, is a known compound (see *J. Heterocycl. Chem.* 1987, 24(5), 1313-16). A convenient preparation of 74 from the 2-aminopyridine of Formula 73 involves protection, ortho-metallation, chlorination and subsequent deprotection. The remaining synthesis is conducted according to the general synthesis described in Scheme 25.

Scheme 27



It is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Steps in the following Examples illustrate a procedure for each step in an overall synthetic transformation, and the starting material for each step may not have necessarily been prepared by a particular preparative run whose procedure is described in other Examples or Steps. Percentages are by weight except for chromatographic solvent mixtures or where otherwise indicated. Parts and percentages for chromatographic solvent mixtures are by

volume unless otherwise indicated. ^1H NMR spectra are reported in ppm downfield from tetramethylsilane; "s" means singlet, "d" means doublet, "t" means triplet, "q" means quartet, "m" means multiplet, "dd" means doublet of doublets, "dt" means doublet of triplets, and "br s" means broad singlet.

5

EXAMPLE 1

Preparation of 2-[3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-6-chloro-8-methyl-4*H*-3,1-benzoxazin-4-one

Step A: Preparation of 2-amino-3-methyl-5-chlorobenzoic acid

To a solution of 2-amino-3-methylbenzoic acid (Aldrich, 15.0 g, 99.2 mmol) in *N,N*-dimethylformamide (50 mL) was added *N*-chlorosuccinimide (13.3 g, 99.2 mmol) and the reaction mixture was heated to 100 °C for 30 minutes. The heat was removed and the reaction mixture was cooled to room temperature and allowed to stand overnight. The reaction mixture was then slowly poured into ice water (250 mL) to precipitate a white solid. The solid was filtered and washed four times with water and then taken up in ethyl acetate (900 mL). The ethyl acetate solution was dried over magnesium sulfate, evaporated under reduced pressure and the residual solid was washed with ether to afford the desired intermediate as a white solid (13.9 g).

^1H NMR (DMSO- d_6) δ 2.11 (s, 3H), 7.22 (s, 1H), 7.55 (s, 1H).

Step B: Preparation of 3-bromo-*N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide

To a solution of *N*-dimethylsulfamoylpyrazole (44.0 g, 0.251 mol) in dry tetrahydrofuran (500 mL) at -78 °C was added dropwise a solution of *n*-butyllithium (2.5 M in hexane, 105.5 mL, 0.264 mol) while maintaining the temperature below -60 °C. A thick solid formed during the addition. Upon completion of the addition the reaction mixture was maintained for an additional 15 minutes, after which time a solution of 1,2-dibromo-tetrachloroethane (90 g, 0.276 mol) in tetrahydrofuran (150 mL) was added dropwise while maintaining the temperature below -70 °C. The reaction mixture turned a clear orange; stirring was continued for an additional 15 minutes. The -78 °C bath was removed and the reaction was quenched with water (600 mL). The reaction mixture was extracted with dichloromethane (4x), and the organic extracts were dried over magnesium sulfate and concentrated. The crude product was further purified by chromatography on silica gel using dichloromethane-hexane (50:50) as eluent to afford the title product as a clear colorless oil (57.04 g).

^1H NMR (CDCl_3) δ 3.07 (d, 6H), 6.44 (m, 1H), 7.62 (m, 1H).

Step C: Preparation of 3-bromopyrazole

To trifluoroacetic acid (70 mL) was slowly added 3-bromo-*N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide (i.e. the bromopyrazole product of Step B) (57.04 g). The reaction

mixture was stirred at room temperature for 30 minutes and then concentrated at reduced pressure. The residue was taken up in hexane, insoluble solids were filtered off, and the hexane was evaporated to afford the crude product as an oil. The crude product was further purified by chromatography on silica gel using ethyl acetate/dichloromethane (10:90) as eluent to afford an oil. The oil was taken up in dichloromethane, neutralized with aqueous sodium bicarbonate solution, extracted with dichloromethane (3x), dried over magnesium sulfate and concentrated to afford the title product as a white solid (25.9 g), m.p. 61–64 °C. ^1H NMR (CDCl_3) δ 6.37 (d, 1H), 7.59 (d, 1H), 12.4 (br s, 1H).

Step D: Preparation of 2-(3-bromo-1H-pyrazol-1-yl)-3-chloropyridine

To a mixture of 2,3-dichloropyridine (27.4 g, 185 mmol) and 3-bromopyrazole (i.e. the product of Step C) (25.4 g, 176 mmol) in dry *N,N*-dimethylformamide (88 mL) was added potassium carbonate (48.6 g, 352 mmol), and the reaction mixture was heated to 125 °C for 18 hours. The reaction mixture was cooled to room temperature and poured into ice water (800 mL). A precipitate formed. The precipitated solids were stirred for 1.5 hours, filtered and washed with water (2x100 mL). The solid filter cake was taken up in dichloromethane and washed sequentially with water, 1N hydrochloric acid, saturated aqueous sodium bicarbonate solution, and brine. The organic extracts were then dried over magnesium sulfate and concentrated to afford 39.9 g of a pink solid. The crude solid was suspended in hexane and stirred vigorously for 1 hour. The solids were filtered, washed with hexane and dried to afford the title product as an off-white powder (30.4 g) determined to be > 94 % pure by NMR. This material was used without further purification in Step D. ^1H NMR (CDCl_3) δ 6.52 (s, 1H), 7.30 (dd, 1H), 7.92 (d, 1H), 8.05 (s, 1H), 8.43 (d, 1H).

Step E: Preparation of 3-bromo-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxylic acid

To a solution of 2-(3-bromo-1H-pyrazol-1-yl)-3-chloropyridine (i.e. the pyrazole product of Step D) (30.4 g, 118 mmol) in dry tetrahydrofuran (250 mL) at –76 °C was added dropwise a solution of lithium diisopropylamide (118 mmol) in tetrahydrofuran at such a rate as to maintain the temperature below –71 °C. The reaction mixture was stirred for 15 minutes at –76 °C, and carbon dioxide was then bubbled through for 10 minutes, causing warming to –57 °C. The reaction mixture was warmed to –20 °C and quenched with water. The reaction mixture was concentrated and then taken up in water (1 L) and ether (500 mL), and then aqueous sodium hydroxide solution (1 N, 20 mL) was added. The phases were separated and the aqueous phase was washed with ether and acidified with hydrochloric acid. The precipitated solids were filtered, washed with water and dried to afford the title product as a tan solid (27.7 g). (Product from another run following similar procedure melted at 200–201 °C.)

^1H NMR ($\text{DMSO}-d_6$) δ 7.25 (s, 1H), 7.68 (dd, 1H), 8.24 (d, 1H), 8.56 (d, 1H).

Step F: Preparation of 2-[3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-6-chloro-8-methyl-4*H*-3,1-benzoxazin-4-one

Methanesulfonyl chloride (1.0 mL, 1.5 g, 13 mmol) was dissolved in acetonitrile (10 mL), and the mixture was cooled to -5°C . A solution of 3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxylic acid (i.e. the pyrazolecarboxylic acid product of Step E) (3.02 g, 10 mmol) and pyridine (1.4 mL, 1.4 g, 17 mmol) in acetonitrile (10 mL) was added dropwise over five minutes at -5 to 0°C . A slurry formed during the addition. The mixture was stirred five minutes at this temperature, and then a mixture of 2-amino-3-methyl-5-chlorobenzoic acid (i.e. the product of Step A) (1.86 g, 10 mmol) and pyridine (2.8 mL, 2.7 g, 35 mmol) in acetonitrile (10 mL) was added, rinsing with more acetonitrile (5 mL). The mixture was stirred 15 minutes at -5 to 0°C , and then methanesulfonyl chloride (1.0 mL, 1.5 mL, 13 mmol) in acetonitrile (5 mL) was added dropwise over five minutes at a temperature of -5 to 0°C . The reaction mixture was stirred 15 minutes more at this temperature, then allowed to warm slowly to room temperature, and stirred 4 hours. Water (20 mL) was added dropwise, and the mixture was stirred 15 minutes. The mixture was filtered, and the solids were washed with 2:1 acetonitrile–water (3×3 mL), then with acetonitrile (2×3 mL), and dried under nitrogen to afford the title product as a light yellow powder, 4.07 g (90.2% crude yield), melting at 203 – 205°C . HPLC of the product using a Zorbax® RX-C8 chromatography column (4.6 mm \times 25 cm, eluent 25–95% acetonitrile/ pH 3 water) showed a major peak corresponding to the title compound and having 95.7% of total chromatogram peak area.

^1H NMR (DMSO- d_6) δ 1.72 (s, 3H) 7.52 (s, 1H), 7.72–7.78 (m, 2H), 7.88 (m, 1H), 8.37 (dd, 1H), 8.62 (dd, 1H).

Preparation of 2-[3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-6-chloro-8-methyl-4*H*-3,1-benzoxazin-4-one in Alternate Solvents

By the method described for Example 1, Step F, except that the solvents shown in Table 1 were used in place of acetonitrile. In those cases where the filtrate formed two phases, the organic phase was separated, dried over magnesium sulfate, filtered, and evaporated, and the residue was triturated with acetonitrile to afford a second crop of 2-[3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-6-chloro-8-methyl-4*H*-3,1-benzoxazin-4-one, though of lower purity.

Table 1

Example	Solvent	First Crop	Second Crop	Total % Yield
		(% Yield)	(% Yield)	
2	Ethyl Acetate	62.2	21	83.3
3	Acetone	86.2	—	86.2
4	Tetrahydrofuran	58.9	21	80.0

<u>Example</u>	<u>Solvent</u>	<u>First Crop</u> <u>(% Yield)</u>	<u>Second Crop</u> <u>(% Yield)</u>	<u>Total % Yield</u>
5	Dichloromethane	79.5	10	89.7

Preparation of 2-[3-bromo-1-(3-chloro-2-pyridinyl)-1H-pyrazol-5-yl]-6-chloro-8-methyl-4H-3,1-benzoxazin-4-one with Alternate Bases

By the method described for Example 1, Step F, except that the bases shown in Table 2 were used in place of pyridine. In each case, the molar quantity of the base used was the same as the molar quantity of pyridine for which it was substituted.

Table 2

<u>Example</u>	<u>Base</u>	<u>% Yield</u>	<u>HPLC Area %</u>
6	3-picoline	92.4	95.2
7	2,6-lutidine	95.5	92.1
8	Collidine	90.4	78.1
9	2-picoline	93.3	90.2
10	4-picoline	68.5	78.0

Preparation of 2-[3-bromo-1-(3-chloro-2-pyridinyl)-1H-pyrazol-5-yl]-6-chloro-8-methyl-4H-3,1-benzoxazin-4-one with Alternate Sulfonyl Chlorides

By the method described for Example 1, Step F, except that the sulfonyl chlorides shown in Table 3 were used in place of methanesulfonyl chloride. In each case, the molar quantity of the sulfonyl chloride used was the same as the molar quantity of methanesulfonyl chloride for which it was substituted.

Table 3

<u>Example</u>	<u>Sulfonyl Chloride</u>	<u>% Yield</u>	<u>HPLC Area %</u>
11	Benzenesulfonyl Chloride	87.5	96.5
12	<i>n</i> -Propanesulfonyl Chloride	93.5	95.0

EXAMPLE 13

Preparation of 6-chloro-2-[3-chloro-1-(3-chloro-2-pyridinyl)-1H-pyrazol-5-yl]-8-methyl-4H-3,1-benzoxazin-4-one

Step A: Preparation of 3-chloro-*N,N*-dimethyl-1H-pyrazole-1-sulfonamide

To a solution of *N*-dimethylsulfamoylpyrazole (188.0 g, 1.07 mol) in dry tetrahydrofuran (1500 mL) at -78°C was added dropwise a solution of 2.5 M *n*-butyllithium (472 mL, 1.18 mol) in hexane while maintaining the temperature below -65°C . Upon completion of the addition the reaction mixture was maintained at -78°C for an additional 45 minutes, after which time a solution of hexachloroethane (279 g, 1.18 mol)

in tetrahydrofuran (120 mL) was added dropwise. The reaction mixture was maintained for an hour at -78°C , warmed to -20°C and then quenched with water (1 L). The reaction mixture was extracted with dichloromethane (4 x 500 mL); the organic extracts were dried over magnesium sulfate and concentrated. The crude product was further purified by chromatography on silica gel using dichloromethane as eluent to afford the title compound as a yellow oil (160 g).

$^1\text{H NMR}$ (CDCl_3) δ 3.07 (d, 6H), 6.33 (s, 1H), 7.61 (s, 1H).

Step B: Preparation of 3-chloropyrazole

To trifluoroacetic acid (290 mL) was added dropwise 3-chloro-*N,N*-dimethyl-1*H*-pyrazole-1-sulfonamide (i.e. the chloropyrazole product of Step A) (160 g), and the reaction mixture was stirred at room temperature for 1.5 hours and then concentrated at reduced pressure. The residue was taken up in hexane, insoluble solids were filtered off, and the hexane was concentrated to afford the crude product as an oil. The crude product was further purified by chromatography on silica gel using ether/hexane (40:60) as eluent to afford the title compound as a yellow oil (64.44 g).

$^1\text{H NMR}$ (CDCl_3) δ 6.39 (s, 1H), 7.66 (s, 1H), 9.6 (br s, 1H).

Step C: Preparation of 3-chloro-2-(3-chloro-1*H*-pyrazol-1-yl)pyridine

To a mixture of 2,3-dichloropyridine (92.60 g, 0.629 mol) and 3-chloropyrazole (i.e. the product of Step B) (64.44 g, 0.629 mol) in *N,N*-dimethylformamide (400 mL) was added potassium carbonate (147.78 g, 1.06 mol), and the reaction mixture was then heated to 100°C for 36 hours. The reaction mixture was cooled to room temperature and slowly poured into ice water. The precipitated solids were filtered and washed with water. The solid filter cake was taken up in ethyl acetate, dried over magnesium sulfate and concentrated. The crude solid was chromatographed on silica gel using 20% ethyl acetate/hexane as eluent to afford the title product as a white solid (39.75 g).

$^1\text{H NMR}$ (CDCl_3) δ 6.43 (s, 1H), 7.26 (m, 1H), 7.90 (d, 1H), 8.09 (s, 1H), 8.41 (d, 1H).

Step D: Preparation of 3-chloro-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxylic acid

To a solution of 3-chloro-2-(3-chloro-1*H*-pyrazol-1-yl)pyridine (i.e. the pyrazole product of Step C) (39.75 g, 186 mmol) in dry tetrahydrofuran (400 mL) at -78°C was added dropwise a solution of 2.0 M lithium diisopropylamide (93 mL, 186 mmol) in tetrahydrofuran. Carbon dioxide was bubbled through the amber solution for 14 minutes, after which time the solution became pale brownish-yellow. The reaction was made basic with 1 N aqueous sodium hydroxide solution and extracted with ether (2 x 500 mL). The aqueous extracts were acidified with 6 N hydrochloric acid and extracted with ethyl acetate (3 x 500 mL). The ethyl acetate extracts were dried over magnesium sulfate and

concentrated to afford the title product as an off-white solid (42.96 g). (Product from another run following a similar procedure melted at 198–199 °C.)

¹H NMR (DMSO-*d*₆) δ 6.99 (s, 1H), 7.45 (m, 1H), 7.93 (d, 1H), 8.51 (d, 1H).

Step E: Preparation of 6-chloro-2-[3-chloro-1-(3-chloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-8-methyl-4*H*-3,1-benzoxazin-4-one

Methanesulfonyl chloride (1.0 mL, 1.5 g, 13 mmol) was dissolved in acetonitrile (10 mL), and the mixture was cooled to –5 °C. A solution of 3-chloro-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxylic acid (i.e. the carboxylic acid product of Step D) (2.58 g, 10 mmol) and pyridine (1.4 mL, 1.4 g, 17 mmol) in acetonitrile (10 mL) was added dropwise over five minutes at –5 to 0 °C. A slurry formed during the addition. The mixture was stirred five minutes at this temperature, and then 2-amino-3-methyl-5-chlorobenzoic acid (i.e. the product from Example 1, Step A) (1.86 g, 10 mmol) was added all at once. Then a solution of pyridine (2.8 mL, 2.7 g, 35 mmol) in acetonitrile (10 mL) was added dropwise over five minutes at –5 to 0 °C. The mixture was stirred 15 minutes at –5 to 0 °C, and then methanesulfonyl chloride (1.0 mL, 1.5 mL, 13 mmol) in acetonitrile (5 mL) was added dropwise over five minutes at –5 to 0 °C. The reaction mixture was stirred 15 minutes at this temperature, then allowed to warm slowly to room temperature, and stirred for 4 hours. Water (15 mL) was added dropwise, and the mixture was stirred 15 minutes. Then the mixture was filtered, and the solids were washed with 2:1 acetonitrile–water (3 × 3 mL), then with acetonitrile (2 × 3 mL), and dried under nitrogen to afford the title product as a pale yellow powder, 3.83 g (94.0% crude yield), melting at 199–201 °C. HPLC of the product using a Zorbax® RX-C8 chromatography column (4.6 mm × 25 cm, eluent 25–95% acetonitrile/ pH 3 water) showed a major peak corresponding to the title compound and having 97.8% of total chromatogram peak area.

¹H NMR (DMSO-*d*₆) δ 1.72 (s, 3H), 7.48 (s, 1H), 7.74–7.80 (m, 2H), 7.87 (m, 1H), 8.37 (dd, 1H), 8.62 (dd, 1H).

EXAMPLE 14

Preparation of a mixture of 2-[3-bromo-1-(3,4-dichloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-6-chloro-8-methyl-4*H*-3,1-benzoxazin-4-one and 2-[3-bromo-1-(3,6-dichloro-2-pyridinyl)-1*H*-pyrazol-5-yl]-6-chloro-8-methyl-4*H*-3,1-benzoxazin-4-one

By the method described above for preparation of Example 13, except that a mixture of 3-bromo-1-(3,4-dichloro-2-pyridinyl)-1*H*-pyrazole-5-carboxylic acid and 3-bromo-1-(3,6-dichloro-2-pyridinyl)-1*H*-pyrazole-5-carboxylic acid (mixture of isomers, 3.37 g, 10 mmol) was used in place of 3-chloro-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxylic acid. The title compound mixture was obtained as light yellow solids, 4.35 g (89.4% crude yield), m. p. 195–210°C. HPLC of the product on a Zorbax RX-C8 column (4.6 mm × 25 cm, eluent 25–95% acetonitrile/ pH 3 water) showed 86.9 area % for the mixture of isomers.


^1H NMR ($\text{DMSO}-d_6$) δ 1.75 and 1.80 (each s, total 3H), 7.56 and 7.56 (each s, total 1H), 7.80 and 7.90 (each m, total 2H), 7.96 and 8.12 (each d, total 1H), 8.44 and 8.61 (each d, total 1H).

Preparation of 2-[3-bromo-1-(3-chloro-1-oxido-2-pyridinyl)-1H-pyrazol-5-yl]-6-chloro-8-methyl-4H-3,1-benzoxazin-4-one

Methanesulfonyl chloride (0.50 mL, 0.74 g, 6.5 mmol) was dissolved in acetonitrile (5 mL) and the mixture was cooled to -5°C . A mixture of 3-chloro-1-(3-chloro-1-oxido-2-pyridinyl)-1H-pyrazole-5-carboxylic acid (1.59 g, 5.0 mmol) and pyridine (0.69 mL, 0.67 g, 8.5 mmol) in acetonitrile (5 mL) formed a slurry and was added dropwise over five minutes at -5 to 0°C . The mixture was stirred five minutes at this temperature, then 2-amino-3-methyl-5-chlorobenzoic acid (i.e. the product from Example 1, Step A) (0.93 g, 5.0 mmol) was added all at once. Then a solution of pyridine (1.4 mL, 1.4 g, 17 mmol) in acetonitrile (2.5 mL) was added dropwise over five minutes at -5 to 0°C . The mixture was stirred 15 minutes at -5 to 0°C , then methanesulfonyl chloride (0.50 mL, 0.74 g, 6.5 mmol) in acetonitrile (2.5 mL) was added dropwise over five minutes at -5 to 0°C . The reaction mixture was stirred 15 minutes at this temperature, then allowed to warm slowly to room temperature, and stirred 3 hours. Water (7.5 mL) was added dropwise, and the mixture was stirred 15 minutes. Then the mixture was filtered and the solids were washed with 2:1 acetonitrile:water (3×1.5 mL), then with acetonitrile (2×1.5 mL), and dried under nitrogen to afford the title compound as an off-white powder, 2.05 g (87.6% crude yield), m. p. $240-245^\circ\text{C}$ (decomposed). HPLC of the product on a Zorbax RX-C8 column (4.6 mm \times 25 cm, eluent 25-95% acetonitrile/ pH 3 water) showed 96.1 area % of the title compound. ^1H NMR ($\text{DMSO}-d_6$) δ 1.87 (s,3H), 7.61 (s,1H), 7.66-7.74 (m,1H), 7.84 (d,1H), 7.89 (d,1H), 8.52 (d,1H).

By the methods and procedures described herein together with methods known in the art, the following compounds of Table 4 can be prepared.

TABLE 4




$\underline{\text{R}^4}$	$\underline{\text{R}^5}$	$\underline{\text{R}^9}$	$\underline{\text{R}^7}$	$\underline{\text{R}^4}$	$\underline{\text{R}^5}$	$\underline{\text{R}^9}$	$\underline{\text{R}^7}$	$\underline{\text{R}^4}$	$\underline{\text{R}^5}$	$\underline{\text{R}^9}$	$\underline{\text{R}^7}$
CH_3	F	CF_3	Cl	Cl	F	CF_3	Cl	CH_3	F	OCH_2CF_3	Cl

R^4	R^5	R^9	R^7	R^4	R^5	R^9	R^7	R^4	R^5	R^9	R^7
CH ₃	F	CF ₃	Br	Cl	F	CF ₃	Br	CH ₃	F	OCH ₂ CF ₃	Br
CH ₃	F	Cl	Cl	Cl	F	Cl	Cl	CH ₃	F	OCHF ₂	Cl
CH ₃	F	Cl	Br	Cl	F	Cl	Br	CH ₃	F	OCHF ₂	Br
CH ₃	F	Br	Cl	Cl	F	Br	Cl	Cl	F	OCH ₂ CF ₃	Cl
CH ₃	F	Br	Br	Cl	F	Br	Br	Cl	F	OCH ₂ CF ₃	Br
CH ₃	Cl	CF ₃	Cl	Cl	Cl	CF ₃	Cl	Cl	F	OCHF ₂	Cl
CH ₃	Cl	CF ₃	Br	Cl	Cl	CF ₃	Br	Cl	F	OCHF ₂	Br
CH ₃	Cl	Cl	Cl	Cl	Cl	Cl	Cl	CH ₃	Cl	OCH ₂ CF ₃	Cl
CH ₃	Cl	Cl	Br	Cl	Cl	Cl	Br	CH ₃	Cl	OCH ₂ CF ₃	Br
CH ₃	Cl	Br	Cl	Cl	Cl	Br	Cl	CH ₃	Cl	OCHF ₂	Cl
CH ₃	Cl	Br	Br	Cl	Cl	Br	Br	CH ₃	Cl	OCHF ₂	Br
CH ₃	Br	CF ₃	Cl	Cl	Br	CF ₃	Cl	Cl	Cl	OCH ₂ CF ₃	Cl
CH ₃	Br	CF ₃	Br	Cl	Br	CF ₃	Br	Cl	Cl	OCH ₂ CF ₃	Br
CH ₃	Br	Cl	Cl	Cl	Br	Cl	Cl	Cl	Cl	OCHF ₂	Cl
CH ₃	Br	Cl	Br	Cl	Br	Cl	Br	Cl	Cl	OCHF ₂	Br
CH ₃	Br	Br	Cl	Cl	Br	Br	Cl	CH ₃	Br	OCH ₂ CF ₃	Cl
CH ₃	Br	Br	Br	Cl	Br	Br	Br	CH ₃	Br	OCH ₂ CF ₃	Br
CH ₃	I	CF ₃	Cl	Cl	I	CF ₃	Cl	CH ₃	Br	OCHF ₂	Cl
CH ₃	I	CF ₃	Br	Cl	I	CF ₃	Br	CH ₃	Br	OCHF ₂	Br
CH ₃	I	Cl	Cl	Cl	I	Cl	Cl	Cl	Br	OCH ₂ CF ₃	Cl
CH ₃	I	Cl	Br	Cl	I	Cl	Br	Cl	Br	OCH ₂ CF ₃	Br
CH ₃	I	Br	Cl	Cl	I	Br	Cl	Cl	Br	OCHF ₂	Cl
CH ₃	I	Br	Br	Cl	I	Br	Br	Cl	Br	OCHF ₂	Br
CH ₃	CF ₃	CF ₃	Cl	Cl	CF ₃	CF ₃	Cl	Cl	I	OCH ₂ CF ₃	Cl
CH ₃	CF ₃	CF ₃	Br	Cl	CF ₃	CF ₃	Br	Cl	I	OCH ₂ CF ₃	Br
CH ₃	CF ₃	Cl	Cl	Cl	CF ₃	Cl	Cl	Cl	I	OCHF ₂	Cl
CH ₃	CF ₃	Cl	Br	Cl	CF ₃	Cl	Br	Cl	I	OCHF ₂	Br
CH ₃	CF ₃	Br	Cl	Cl	CF ₃	Br	Cl	CH ₃	I	OCH ₂ CF ₃	Cl
CH ₃	CF ₃	Br	Br	Cl	CF ₃	Br	Br	CH ₃	I	OCH ₂ CF ₃	Br
CH ₃	Cl	Cl	Cl	Cl	Cl	Cl	Cl	CH ₃	I	OCHF ₂	Cl
F	F	CF ₃	Cl	Br	F	CF ₃	Cl	CH ₃	I	OCHF ₂	Br
F	F	CF ₃	Br	Br	F	CF ₃	Br	Cl	CF ₃	OCH ₂ CF ₃	Cl
F	F	Cl	Cl	Br	F	Cl	Cl	Cl	CF ₃	OCH ₂ CF ₃	Br
F	F	Cl	Br	Br	F	Cl	Br	Cl	CF ₃	OCHF ₂	Cl
F	F	Br	Cl	Br	F	Br	Cl	Cl	CF ₃	OCHF ₂	Br
F	F	Br	Br	Br	F	Br	Br	CH ₃	CF ₃	OCH ₂ CF ₃	Cl
F	Cl	CF ₃	Cl	Br	Cl	CF ₃	Cl	CH ₃	CF ₃	OCH ₂ CF ₃	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R⁷</u>
F	Cl	CF ₃	Br	Br	Cl	CF ₃	Br	CH ₃	CF ₃	OCHF ₂	Cl
F	Cl	Cl	Cl	Br	Cl	Cl	Cl	CH ₃	CF ₃	OCHF ₂	Br
F	Cl	Cl	Br	Br	Cl	Cl	Br	Br	F	OCH ₂ CF ₃	Cl
F	Cl	Br	Cl	Br	Cl	Br	Cl	Br	F	OCH ₂ CF ₃	Br
F	Cl	Br	Br	Br	Cl	Br	Br	F	F	OCH ₂ CF ₃	Cl
F	Br	CF ₃	Cl	Br	Br	CF ₃	Cl	F	F	OCH ₂ CF ₃	Br
F	Br	CF ₃	Br	Br	Br	CF ₃	Br	Br	Cl	OCH ₂ CF ₃	Cl
F	Br	Cl	Cl	Br	Br	Cl	Cl	Br	Cl	OCH ₂ CF ₃	Br
F	Br	Cl	Br	Br	Br	Cl	Br	F	Cl	OCH ₂ CF ₃	Cl
F	Br	Br	Cl	Br	Br	Br	Cl	F	Cl	OCH ₂ CF ₃	Br
F	Br	Br	Br	Br	Br	Br	Br	Br	Br	OCH ₂ CF ₃	Cl
F	I	CF ₃	Cl	Br	I	CF ₃	Cl	Br	Br	OCH ₂ CF ₃	Br
F	I	CF ₃	Br	Br	I	CF ₃	Br	F	Br	OCH ₂ CF ₃	Cl
F	I	Cl	Cl	Br	I	Cl	Cl	F	Br	OCH ₂ CF ₃	Br
F	I	Cl	Br	Br	I	Cl	Br	F	Br	OCHF ₂	Cl
F	I	Br	Cl	Br	I	Br	Cl	F	Br	OCHF ₂	Br
F	I	Br	Br	Br	I	Br	Br	Br	I	OCH ₂ CF ₃	Cl
F	CF ₃	CF ₃	Cl	Br	CF ₃	CF ₃	Cl	Br	I	OCH ₂ CF ₃	Br
F	CF ₃	CF ₃	Br	Br	CF ₃	CF ₃	Br	Br	I	OCHF ₂	Cl
F	CF ₃	Cl	Cl	Br	CF ₃	Cl	Cl	Br	I	OCHF ₂	Br
F	CF ₃	Cl	Br	Br	CF ₃	Cl	Br	F	I	OCH ₂ CF ₃	Cl
F	CF ₃	Br	Cl	Br	CF ₃	Br	Cl	F	I	OCH ₂ CF ₃	Br
F	CF ₃	Br	Br	Br	CF ₃	Br	Br	Br	CF ₃	OCH ₂ CF ₃	Cl
F	F	OCHF ₂	Cl	Br	F	OCHF ₂	Cl	Br	CF ₃	OCH ₂ CF ₃	Br
F	F	OCHF ₂	Br	Br	F	OCHF ₂	Br	Br	CF ₃	OCHF ₂	Cl
F	Cl	OCHF ₂	Cl	Br	Cl	OCHF ₂	Cl	Br	CF ₃	OCHF ₂	Br
F	Cl	OCHF ₂	Br	Br	Cl	OCHF ₂	Br	F	CF ₃	OCH ₂ CF ₃	Cl
F	I	OCHF ₂	Cl	Br	Br	OCHF ₂	Cl	F	CF ₃	OCH ₂ CF ₃	Br
F	I	OCHF ₂	Br	Br	Br	OCHF ₂	Br	F	CF ₃	OCHF ₂	Cl
								F	CF ₃	OCHF ₂	Br


The fused oxazinone preparation method of the invention can be used to prepare a wide variety of compounds of Formula 1 that are useful as intermediates for the preparation of crop protection agents, pharmaceuticals and other fine chemicals.

Compounds of Formula 1 can be used to prepare compounds of Formula II by reaction
 5 with nucleophiles, optionally in the presence of additional base.



As shown in Scheme 28, reaction of Formula 1 with nucleophiles of Formula 77 wherein R^d can be an optionally substituted carbon moiety (i.e. alcohols) leads to esters of Formula IIa. Reaction of Formula 1 with nucleophiles of Formula 78 wherein R^e and R^f can be independently H or an optionally substituted carbon moiety (i.e. ammonia, primary amines or secondary amines) leads to amides of Formula IIb.


Scheme 28



Typical procedures for preparation of compounds of Formula IIb involve combination of an amine of Formula 78 with the fused oxazinone of Formula 1. The reaction can be run neat or in a variety of suitable solvents including tetrahydrofuran, diethyl ether, dichloromethane or chloroform with optimum temperatures ranging from room temperature to the reflux temperature of the solvent. For references to the chemistry of heterocyclic fused oxazinones see Jakobsen et al, *Biorganic and Medicinal Chemistry*, 2000, 8, 2803-2812 and references cited therein.

As a particular example, compounds of Formula Ia are useful for preparing compounds of Formula III

51



wherein

X is N or CR⁶;

Y is N or CH;

5 R¹ is H;

R² is H or CH₃;

R³ is C₁-C₆ alkyl;

R⁴ is C₁-C₄ alkyl or halogen;

R⁵ is H, C₁-C₄ alkyl, C₁-C₄ haloalkyl or halogen;

10 R⁶ and R⁷ are independently H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, halogen, CN or C₁-C₄ haloalkoxy;

R⁸ is H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, C₃-C₆cycloalkylamino, (C₁-C₄ alkyl)(C₃-C₆cycloalkyl)amino, C₂-C₄ alkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₂-C₆ alkylaminocarbonyl, C₃-C₈ dialkylaminocarbonyl or C₃-C₆ trialkylsilyl; and


R⁹ is CF₃, OCF₃, OCHF₂, OCH₂CF₃, S(O)_pCF₃, S(O)_pCHF₂ or halogen.

20 Compounds of Formula **III** are useful as insecticides, as described, for example, in PCT Publication No. WO 01/70671, published September 27, 2001, as well as in U.S. Patent Application 60/324,173, filed September 21, 2001, U.S. Patent Application 60/323,941, filed September 21, 2001 and U.S. Patent Application 60/369,661, filed April 2, 2002. The preparation of compounds of Formula 1a and Formula **III** is described in U.S. Patent Application _____[BA9307 US PRV], filed concurrently herewith and hereby

25 incorporated herein in its entirety by reference, as well as in U.S. Patent Application 60/369,659, filed April 2, 2002 and U.S. Patent Application 60/369,660, filed April 2, 2002.

30 Compounds of Formula **III** can be prepared by the reaction of benzoxazinones of Formula 1a with C₁-C₆ alkylamines and (C₁-C₆ alkyl)(methyl)amines of Formula 79 as outlined in Scheme 8.

Scheme 29




The reaction can be run neat or in a variety of suitable solvents including tetrahydrofuran, diethyl ether, dichloromethane or chloroform with optimum temperatures ranging from room temperature to the reflux temperature of the solvent. The general reaction of benzoxazinones with amines to produce anthranilamides is well documented in the chemical literature. For a review of benzoxazinone chemistry see Jakobsen et al., *Biorganic and Medicinal Chemistry* 2000, 8, 2095–2103 and references cited within. See also Coppola, *J. Heterocyclic Chemistry* 1999, 36, 563–588.

- Preferred are methods for preparing compounds of Formula 1a or Formula **III** wherein
- X is N;
 - Y is N;
 - R² is H or CH₃;
 - R³ is C₁–C₄ alkyl;
 - R⁴ is CH₃, F, Cl or Br;
 - R⁵ is CF₃, F, Cl, Br or I;
 - R⁷ is Cl or Br;
 - R⁸ is H, and
 - R⁹ is CF₃, OCHF₂, OCH₂CF₃, Cl or Br.
- Also of particular note are compounds of Formula **III** wherein R⁹ is OCHF₂.

By the procedures described herein together with methods known in the art, the following compounds of Table 5 can be prepared. The following abbreviations are used in the Tables which follow: *t* means tertiary, *s* means secondary, *n* means normal, *i* means iso,

Me means methyl, Et means ethyl, Pr means propyl, *i*-Pr means isopropyl, and Bu means butyl.

TABLE 5



<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	F	CF ₃	Me	H	Cl	Cl	F	CF ₃	Me	H	Cl
CH ₃	F	CF ₃	Et	H	Cl	Cl	F	CF ₃	Et	H	Cl
CH ₃	F	CF ₃	<i>i</i> -Pr	H	Cl	Cl	F	CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	F	CF ₃	<i>t</i> -Bu	H	Cl	Cl	F	CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	F	CF ₃	Me	Me	Cl	Cl	F	CF ₃	Me	Me	Cl
CH ₃	F	CF ₃	Me	H	Br	Cl	F	CF ₃	Me	H	Br
CH ₃	F	CF ₃	Et	H	Br	Cl	F	CF ₃	Et	H	Br
CH ₃	F	CF ₃	<i>i</i> -Pr	H	Br	Cl	F	CF ₃	<i>i</i> -Pr	H	Br
CH ₃	F	CF ₃	<i>t</i> -Bu	H	Br	Cl	F	CF ₃	<i>t</i> -Bu	H	Br
CH ₃	F	CF ₃	Me	Me	Br	Cl	F	CF ₃	Me	Me	Br
CH ₃	F	Cl	Me	H	Cl	Cl	F	Cl	Me	H	Cl
CH ₃	F	Cl	Et	H	Cl	Cl	F	Cl	Et	H	Cl
CH ₃	F	Cl	<i>i</i> -Pr	H	Cl	Cl	F	Cl	<i>i</i> -Pr	H	Cl
CH ₃	F	Cl	<i>t</i> -Bu	H	Cl	Cl	F	Cl	<i>t</i> -Bu	H	Cl
CH ₃	F	Cl	Me	Me	Cl	Cl	F	Cl	Me	Me	Cl
CH ₃	F	Cl	Me	H	Br	Cl	F	Cl	Me	H	Br
CH ₃	F	Cl	Et	H	Br	Cl	F	Cl	Et	H	Br
CH ₃	F	Cl	<i>i</i> -Pr	H	Br	Cl	F	Cl	<i>i</i> -Pr	H	Br
CH ₃	F	Cl	<i>t</i> -Bu	H	Br	Cl	F	Cl	<i>t</i> -Bu	H	Br
CH ₃	F	Cl	Me	Me	Br	Cl	F	Cl	Me	Me	Br
CH ₃	F	Br	Me	H	Cl	Cl	F	Br	Me	H	Cl
CH ₃	F	Br	Et	H	Cl	Cl	F	Br	Et	H	Cl
CH ₃	F	Br	<i>i</i> -Pr	H	Cl	Cl	F	Br	<i>i</i> -Pr	H	Cl
CH ₃	F	Br	<i>t</i> -Bu	H	Cl	Cl	F	Br	<i>t</i> -Bu	H	Cl
CH ₃	F	Br	Me	Me	Cl	Cl	F	Br	Me	Me	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	F	Br	Me	H	Br	Cl	F	Br	Me	H	Br
CH ₃	F	Br	Et	H	Br	Cl	F	Br	Et	H	Br
CH ₃	F	Br	<i>i</i> -Pr	H	Br	Cl	F	Br	<i>i</i> -Pr	H	Br
CH ₃	F	Br	<i>t</i> -Bu	H	Br	Cl	F	Br	<i>t</i> -Bu	H	Br
CH ₃	F	Br	Me	Me	Br	Cl	F	Br	Me	Me	Br
CH ₃	F	OCH ₂ CF ₃	Me	H	Cl	Cl	F	OCH ₂ CF ₃	Me	H	Cl
CH ₃	F	OCH ₂ CF ₃	Et	H	Cl	Cl	F	OCH ₂ CF ₃	Et	H	Cl
CH ₃	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Cl	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Cl	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	F	OCH ₂ CF ₃	Me	Me	Cl	Cl	F	OCH ₂ CF ₃	Me	Me	Cl
CH ₃	F	OCH ₂ CF ₃	Me	H	Br	Cl	F	OCH ₂ CF ₃	Me	H	Br
CH ₃	F	OCH ₂ CF ₃	Et	H	Br	Cl	F	OCH ₂ CF ₃	Et	H	Br
CH ₃	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Cl	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
CH ₃	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Cl	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
CH ₃	F	OCH ₂ CF ₃	Me	Me	Br	Cl	F	OCH ₂ CF ₃	Me	Me	Br
CH ₃	Cl	CF ₃	Me	H	Cl	Cl	Cl	CF ₃	Me	H	Cl
CH ₃	Cl	CF ₃	Et	H	Cl	Cl	Cl	CF ₃	Et	H	Cl
CH ₃	Cl	CF ₃	<i>i</i> -Pr	H	Cl	Cl	Cl	CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	Cl	CF ₃	<i>t</i> -Bu	H	Cl	Cl	Cl	CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	Cl	CF ₃	Me	Me	Cl	Cl	Cl	CF ₃	Me	Me	Cl
CH ₃	Cl	CF ₃	Me	H	Br	Cl	Cl	CF ₃	Me	H	Br
CH ₃	Cl	CF ₃	Et	H	Br	Cl	Cl	CF ₃	Et	H	Br
CH ₃	Cl	CF ₃	<i>i</i> -Pr	H	Br	Cl	Cl	CF ₃	<i>i</i> -Pr	H	Br
CH ₃	Cl	CF ₃	<i>t</i> -Bu	H	Br	Cl	Cl	CF ₃	<i>t</i> -Bu	H	Br
CH ₃	Cl	CF ₃	Me	Me	Br	Cl	Cl	CF ₃	Me	Me	Br
CH ₃	Cl	Cl	Me	H	Cl	Cl	Cl	Cl	Me	H	Cl
CH ₃	Cl	Cl	Et	H	Cl	Cl	Cl	Cl	Et	H	Cl
CH ₃	Cl	Cl	<i>i</i> -Pr	H	Cl	Cl	Cl	Cl	<i>i</i> -Pr	H	Cl
CH ₃	Cl	Cl	<i>t</i> -Bu	H	Cl	Cl	Cl	Cl	<i>t</i> -Bu	H	Cl
CH ₃	Cl	Cl	Me	Me	Cl	Cl	Cl	Cl	Me	Me	Cl
CH ₃	Cl	Cl	Me	H	Br	Cl	Cl	Cl	Me	H	Br
CH ₃	Cl	Cl	Et	H	Br	Cl	Cl	Cl	Et	H	Br
CH ₃	Cl	Cl	<i>i</i> -Pr	H	Br	Cl	Cl	Cl	<i>i</i> -Pr	H	Br
CH ₃	Cl	Cl	<i>t</i> -Bu	H	Br	Cl	Cl	Cl	<i>t</i> -Bu	H	Br
CH ₃	Cl	Cl	Me	Me	Br	Cl	Cl	Cl	Me	Me	Br
CH ₃	Cl	Br	Me	H	Cl	Cl	Cl	Br	Me	H	Cl
CH ₃	Cl	Br	Et	H	Cl	Cl	Cl	Br	Et	H	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	Cl	Br	<i>i</i> -Pr	H	Cl	Cl	Cl	Br	<i>i</i> -Pr	H	Cl
CH ₃	Cl	Br	<i>t</i> -Bu	H	Cl	Cl	Cl	Br	<i>t</i> -Bu	H	Cl
CH ₃	Cl	Br	Me	Me	Cl	Cl	Cl	Br	Me	Me	Cl
CH ₃	Cl	Br	Me	H	Br	Cl	Cl	Br	Me	H	Br
CH ₃	Cl	Br	Et	H	Br	Cl	Cl	Br	Et	H	Br
CH ₃	Cl	Br	<i>i</i> -Pr	H	Br	Cl	Cl	Br	<i>i</i> -Pr	H	Br
CH ₃	Cl	Br	<i>t</i> -Bu	H	Br	Cl	Cl	Br	<i>t</i> -Bu	H	Br
CH ₃	Cl	Br	Me	Me	Br	Cl	Cl	Br	Me	Me	Br
CH ₃	Cl	OCH ₂ CF ₃	Me	H	Cl	Cl	Cl	OCH ₂ CF ₃	Me	H	Cl
CH ₃	Cl	OCH ₂ CF ₃	Et	H	Cl	Cl	Cl	OCH ₂ CF ₃	Et	H	Cl
CH ₃	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Cl	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Cl	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	Cl	OCH ₂ CF ₃	Me	Me	Cl	Cl	Cl	OCH ₂ CF ₃	Me	Me	Cl
CH ₃	Cl	OCH ₂ CF ₃	Me	H	Br	Cl	Cl	OCH ₂ CF ₃	Me	H	Br
CH ₃	Cl	OCH ₂ CF ₃	Et	H	Br	Cl	Cl	OCH ₂ CF ₃	Et	H	Br
CH ₃	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Cl	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
CH ₃	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Cl	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
CH ₃	Cl	OCH ₂ CF ₃	Me	Me	Br	Cl	Cl	OCH ₂ CF ₃	Me	Me	Br
CH ₃	Br	CF ₃	Me	H	Cl	Cl	Br	CF ₃	Me	H	Cl
CH ₃	Br	CF ₃	Et	H	Cl	Cl	Br	CF ₃	Et	H	Cl
CH ₃	Br	CF ₃	<i>i</i> -Pr	H	Cl	Cl	Br	CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	Br	CF ₃	<i>t</i> -Bu	H	Cl	Cl	Br	CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	Br	CF ₃	Me	Me	Cl	Cl	Br	CF ₃	Me	Me	Cl
CH ₃	Br	CF ₃	Me	H	Br	Cl	Br	CF ₃	Me	H	Br
CH ₃	Br	CF ₃	Et	H	Br	Cl	Br	CF ₃	Et	H	Br
CH ₃	Br	CF ₃	<i>i</i> -Pr	H	Br	Cl	Br	CF ₃	<i>i</i> -Pr	H	Br
CH ₃	Br	CF ₃	<i>t</i> -Bu	H	Br	Cl	Br	CF ₃	<i>t</i> -Bu	H	Br
CH ₃	Br	CF ₃	Me	Me	Br	Cl	Br	CF ₃	Me	Me	Br
CH ₃	Br	Cl	Me	H	Cl	Cl	Br	Cl	Me	H	Cl
CH ₃	Br	Cl	Et	H	Cl	Cl	Br	Cl	Et	H	Cl
CH ₃	Br	Cl	<i>i</i> -Pr	H	Cl	Cl	Br	Cl	<i>i</i> -Pr	H	Cl
CH ₃	Br	Cl	<i>t</i> -Bu	H	Cl	Cl	Br	Cl	<i>t</i> -Bu	H	Cl
CH ₃	Br	Cl	Me	Me	Cl	Cl	Br	Cl	Me	Me	Cl
CH ₃	Br	Cl	Me	H	Br	Cl	Br	Cl	Me	H	Br
CH ₃	Br	Cl	Et	H	Br	Cl	Br	Cl	Et	H	Br
CH ₃	Br	Cl	<i>i</i> -Pr	H	Br	Cl	Br	Cl	<i>i</i> -Pr	H	Br
CH ₃	Br	Cl	<i>t</i> -Bu	H	Br	Cl	Br	Cl	<i>t</i> -Bu	H	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	Br	Cl	Me	Me	Br	Cl	Br	Cl	Me	Me	Br
CH ₃	Br	Br	Me	H	Cl	Cl	Br	Br	Me	H	Cl
CH ₃	Br	Br	Et	H	Cl	Cl	Br	Br	Et	H	Cl
CH ₃	Br	Br	<i>i</i> -Pr	H	Cl	Cl	Br	Br	<i>i</i> -Pr	H	Cl
CH ₃	Br	Br	<i>t</i> -Bu	H	Cl	Cl	Br	Br	<i>t</i> -Bu	H	Cl
CH ₃	Br	Br	Me	Me	Cl	Cl	Br	Br	Me	Me	Cl
CH ₃	Br	Br	Me	H	Br	Cl	Br	Br	Me	H	Br
CH ₃	Br	Br	Et	H	Br	Cl	Br	Br	Et	H	Br
CH ₃	Br	Br	<i>i</i> -Pr	H	Br	Cl	Br	Br	<i>i</i> -Pr	H	Br
CH ₃	Br	Br	<i>t</i> -Bu	H	Br	Cl	Br	Br	<i>t</i> -Bu	H	Br
CH ₃	Br	Br	Me	Me	Br	Cl	Br	Br	Me	Me	Br
CH ₃	Br	OCH ₂ CF ₃	Me	H	Cl	Cl	Br	OCH ₂ CF ₃	Me	H	Cl
CH ₃	Br	OCH ₂ CF ₃	Et	H	Cl	Cl	Br	OCH ₂ CF ₃	Et	H	Cl
CH ₃	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Cl	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Cl	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	Br	OCH ₂ CF ₃	Me	Me	Cl	Cl	Br	OCH ₂ CF ₃	Me	Me	Cl
CH ₃	Br	OCH ₂ CF ₃	Me	H	Br	Cl	Br	OCH ₂ CF ₃	Me	H	Br
CH ₃	Br	OCH ₂ CF ₃	Et	H	Br	Cl	Br	OCH ₂ CF ₃	Et	H	Br
CH ₃	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Cl	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
CH ₃	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Cl	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
CH ₃	Br	OCH ₂ CF ₃	Me	Me	Br	Cl	Br	OCH ₂ CF ₃	Me	Me	Br
CH ₃	I	CF ₃	Me	H	Cl	Cl	I	CF ₃	Me	H	Cl
CH ₃	I	CF ₃	Et	H	Cl	Cl	I	CF ₃	Et	H	Cl
CH ₃	I	CF ₃	<i>i</i> -Pr	H	Cl	Cl	I	CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	I	CF ₃	<i>t</i> -Bu	H	Cl	Cl	I	CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	I	CF ₃	Me	Me	Cl	Cl	I	CF ₃	Me	Me	Cl
CH ₃	I	CF ₃	Me	H	Br	Cl	I	CF ₃	Me	H	Br
CH ₃	I	CF ₃	Et	H	Br	Cl	I	CF ₃	Et	H	Br
CH ₃	I	CF ₃	<i>i</i> -Pr	H	Br	Cl	I	CF ₃	<i>i</i> -Pr	H	Br
CH ₃	I	CF ₃	<i>t</i> -Bu	H	Br	Cl	I	CF ₃	<i>t</i> -Bu	H	Br
CH ₃	I	CF ₃	Me	Me	Br	Cl	I	CF ₃	Me	Me	Br
CH ₃	I	Cl	Me	H	Cl	Cl	I	Cl	Me	H	Cl
CH ₃	I	Cl	Et	H	Cl	Cl	I	Cl	Et	H	Cl
CH ₃	I	Cl	<i>i</i> -Pr	H	Cl	Cl	I	Cl	<i>i</i> -Pr	H	Cl
CH ₃	I	Cl	<i>t</i> -Bu	H	Cl	Cl	I	Cl	<i>t</i> -Bu	H	Cl
CH ₃	I	Cl	Me	Me	Cl	Cl	I	Cl	Me	Me	Cl
CH ₃	I	Cl	Me	H	Br	Cl	I	Cl	Me	H	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	I	Cl	Et	H	Br	Cl	I	Cl	Et	H	Br
CH ₃	I	Cl	<i>i</i> -Pr	H	Br	Cl	I	Cl	<i>i</i> -Pr	H	Br
CH ₃	I	Cl	<i>t</i> -Bu	H	Br	Cl	I	Cl	<i>t</i> -Bu	H	Br
CH ₃	I	Cl	Me	Me	Br	Cl	I	Cl	Me	Me	Br
CH ₃	I	Br	Me	H	Cl	Cl	I	Br	Me	H	Cl
CH ₃	I	Br	Et	H	Cl	Cl	I	Br	Et	H	Cl
CH ₃	I	Br	<i>i</i> -Pr	H	Cl	Cl	I	Br	<i>i</i> -Pr	H	Cl
CH ₃	I	Br	<i>t</i> -Bu	H	Cl	Cl	I	Br	<i>t</i> -Bu	H	Cl
CH ₃	I	Br	Me	Me	Cl	Cl	I	Br	Me	Me	Cl
CH ₃	I	Br	Me	H	Br	Cl	I	Br	Me	H	Br
CH ₃	I	Br	Et	H	Br	Cl	I	Br	Et	H	Br
CH ₃	I	Br	<i>i</i> -Pr	H	Br	Cl	I	Br	<i>i</i> -Pr	H	Br
CH ₃	I	Br	<i>t</i> -Bu	H	Br	Cl	I	Br	<i>t</i> -Bu	H	Br
CH ₃	I	Br	Me	Me	Br	Cl	I	Br	Me	Me	Br
CH ₃	I	OCH ₂ CF ₃	Me	H	Cl	Cl	I	OCH ₂ CF ₃	Me	H	Cl
CH ₃	I	OCH ₂ CF ₃	Et	H	Cl	Cl	I	OCH ₂ CF ₃	Et	H	Cl
CH ₃	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Cl	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Cl	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	I	OCH ₂ CF ₃	Me	Me	Cl	Cl	I	OCH ₂ CF ₃	Me	Me	Cl
CH ₃	I	OCH ₂ CF ₃	Me	H	Br	Cl	I	OCH ₂ CF ₃	Me	H	Br
CH ₃	I	OCH ₂ CF ₃	Et	H	Br	Cl	I	OCH ₂ CF ₃	Et	H	Br
CH ₃	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Cl	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
CH ₃	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Cl	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
CH ₃	I	OCH ₂ CF ₃	Me	Me	Br	Cl	I	OCH ₂ CF ₃	Me	Me	Br
CH ₃	CF ₃	CF ₃	Me	H	Cl	Cl	CF ₃	CF ₃	Me	H	Cl
CH ₃	CF ₃	CF ₃	Et	H	Cl	Cl	CF ₃	CF ₃	Et	H	Cl
CH ₃	CF ₃	CF ₃	<i>i</i> -Pr	H	Cl	Cl	CF ₃	CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	CF ₃	CF ₃	<i>t</i> -Bu	H	Cl	Cl	CF ₃	CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	CF ₃	CF ₃	Me	Me	Cl	Cl	CF ₃	CF ₃	Me	Me	Cl
CH ₃	CF ₃	CF ₃	Me	H	Br	Cl	CF ₃	CF ₃	Me	H	Br
CH ₃	CF ₃	CF ₃	Et	H	Br	Cl	CF ₃	CF ₃	Et	H	Br
CH ₃	CF ₃	CF ₃	<i>i</i> -Pr	H	Br	Cl	CF ₃	CF ₃	<i>i</i> -Pr	H	Br
CH ₃	CF ₃	CF ₃	<i>t</i> -Bu	H	Br	Cl	CF ₃	CF ₃	<i>t</i> -Bu	H	Br
CH ₃	CF ₃	CF ₃	Me	Me	Br	Cl	CF ₃	CF ₃	Me	Me	Br
CH ₃	CF ₃	Cl	Me	H	Cl	Cl	CF ₃	Cl	Me	H	Cl
CH ₃	CF ₃	Cl	Et	H	Cl	Cl	CF ₃	Cl	Et	H	Cl
CH ₃	CF ₃	Cl	<i>i</i> -Pr	H	Cl	Cl	CF ₃	Cl	<i>i</i> -Pr	H	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	CF ₃	Cl	<i>t</i> -Bu	H	Cl	Cl	CF ₃	Cl	<i>t</i> -Bu	H	Cl
CH ₃	CF ₃	Cl	Me	Me	Cl	Cl	CF ₃	Cl	Me	Me	Cl
CH ₃	CF ₃	Cl	Me	H	Br	Cl	CF ₃	Cl	Me	H	Br
CH ₃	CF ₃	Cl	Et	H	Br	Cl	CF ₃	Cl	Et	H	Br
CH ₃	CF ₃	Cl	<i>i</i> -Pr	H	Br	Cl	CF ₃	Cl	<i>i</i> -Pr	H	Br
CH ₃	CF ₃	Cl	<i>t</i> -Bu	H	Br	Cl	CF ₃	Cl	<i>t</i> -Bu	H	Br
CH ₃	CF ₃	Cl	Me	Me	Br	Cl	CF ₃	Cl	Me	Me	Br
CH ₃	CF ₃	Br	Me	H	Cl	Cl	CF ₃	Br	Me	H	Cl
CH ₃	CF ₃	Br	Et	H	Cl	Cl	CF ₃	Br	Et	H	Cl
CH ₃	CF ₃	Br	<i>i</i> -Pr	H	Cl	Cl	CF ₃	Br	<i>i</i> -Pr	H	Cl
CH ₃	CF ₃	Br	<i>t</i> -Bu	H	Cl	Cl	CF ₃	Br	<i>t</i> -Bu	H	Cl
CH ₃	CF ₃	Br	Me	Me	Cl	Cl	CF ₃	Br	Me	Me	Cl
CH ₃	CF ₃	Br	Me	H	Br	Cl	CF ₃	Br	Me	H	Br
CH ₃	CF ₃	Br	Et	H	Br	Cl	CF ₃	Br	Et	H	Br
CH ₃	CF ₃	Br	<i>i</i> -Pr	H	Br	Cl	CF ₃	Br	<i>i</i> -Pr	H	Br
CH ₃	CF ₃	Br	<i>t</i> -Bu	H	Br	Cl	CF ₃	Br	<i>t</i> -Bu	H	Br
CH ₃	CF ₃	Br	Me	Me	Br	Cl	CF ₃	Br	Me	Me	Br
CH ₃	CF ₃	OCH ₂ CF ₃	Me	H	Cl	Cl	CF ₃	OCH ₂ CF ₃	Me	H	Cl
CH ₃	CF ₃	OCH ₂ CF ₃	Et	H	Cl	Cl	CF ₃	OCH ₂ CF ₃	Et	H	Cl
CH ₃	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Cl	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
CH ₃	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Cl	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
CH ₃	CF ₃	OCH ₂ CF ₃	Me	Me	Cl	Cl	CF ₃	OCH ₂ CF ₃	Me	Me	Cl
CH ₃	CF ₃	OCH ₂ CF ₃	Me	H	Br	Cl	CF ₃	OCH ₂ CF ₃	Me	H	Br
CH ₃	CF ₃	OCH ₂ CF ₃	Et	H	Br	Cl	CF ₃	OCH ₂ CF ₃	Et	H	Br
CH ₃	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Cl	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
CH ₃	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Cl	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
CH ₃	CF ₃	OCH ₂ CF ₃	Me	Me	Br	Cl	CF ₃	OCH ₂ CF ₃	Me	Me	Br
CH ₃	Cl	Cl	<i>n</i> -Pr	H	Cl	Cl	Cl	Cl	<i>n</i> -Pr	H	Cl
CH ₃	Cl	Cl	<i>n</i> -Bu	H	Cl	Cl	Cl	Cl	<i>n</i> -Bu	H	Cl
CH ₃	Cl	Cl	<i>s</i> -Bu	H	Cl	Cl	Cl	Cl	<i>s</i> -Bu	H	Cl
CH ₃	Cl	Cl	<i>i</i> -Bu	H	Cl	Cl	Cl	Cl	<i>i</i> -Bu	H	Cl
CH ₃	Cl	Cl	Et	Me	Cl	Cl	Cl	Cl	Et	Me	Cl
F	F	CF ₃	Me	H	Cl	Br	F	CF ₃	Me	H	Cl
F	F	CF ₃	Et	H	Cl	Br	F	CF ₃	Et	H	Cl
F	F	CF ₃	<i>i</i> -Pr	H	Cl	Br	F	CF ₃	<i>i</i> -Pr	H	Cl
F	F	CF ₃	<i>t</i> -Bu	H	Cl	Br	F	CF ₃	<i>t</i> -Bu	H	Cl
F	F	CF ₃	Me	Me	Cl	Br	F	CF ₃	Me	Me	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	F	CF ₃	Me	H	Br	Br	F	CF ₃	Me	H	Br
F	F	CF ₃	Et	H	Br	Br	F	CF ₃	Et	H	Br
F	F	CF ₃	<i>i</i> -Pr	H	Br	Br	F	CF ₃	<i>i</i> -Pr	H	Br
F	F	CF ₃	<i>t</i> -Bu	H	Br	Br	F	CF ₃	<i>t</i> -Bu	H	Br
F	F	CF ₃	Me	Me	Br	Br	F	CF ₃	Me	Me	Br
F	F	Cl	Me	H	Cl	Br	F	Cl	Me	H	Cl
F	F	Cl	Et	H	Cl	Br	F	Cl	Et	H	Cl
F	F	Cl	<i>i</i> -Pr	H	Cl	Br	F	Cl	<i>i</i> -Pr	H	Cl
F	F	Cl	<i>t</i> -Bu	H	Cl	Br	F	Cl	<i>t</i> -Bu	H	Cl
F	F	Cl	Me	Me	Cl	Br	F	Cl	Me	Me	Cl
F	F	Cl	Me	H	Br	Br	F	Cl	Me	H	Br
F	F	Cl	Et	H	Br	Br	F	Cl	Et	H	Br
F	F	Cl	<i>i</i> -Pr	H	Br	Br	F	Cl	<i>i</i> -Pr	H	Br
F	F	Cl	<i>t</i> -Bu	H	Br	Br	F	Cl	<i>t</i> -Bu	H	Br
F	F	Cl	Me	Me	Br	Br	F	Cl	Me	Me	Br
F	F	Br	Me	H	Cl	Br	F	Br	Me	H	Cl
F	F	Br	Et	H	Cl	Br	F	Br	Et	H	Cl
F	F	Br	<i>i</i> -Pr	H	Cl	Br	F	Br	<i>i</i> -Pr	H	Cl
F	F	Br	<i>t</i> -Bu	H	Cl	Br	F	Br	<i>t</i> -Bu	H	Cl
F	F	Br	Me	Me	Cl	Br	F	Br	Me	Me	Cl
F	F	Br	Me	H	Br	Br	F	Br	Me	H	Br
F	F	Br	Et	H	Br	Br	F	Br	Et	H	Br
F	F	Br	<i>i</i> -Pr	H	Br	Br	F	Br	<i>i</i> -Pr	H	Br
F	F	Br	<i>t</i> -Bu	H	Br	Br	F	Br	<i>t</i> -Bu	H	Br
F	F	Br	Me	Me	Br	Br	F	Br	Me	Me	Br
F	F	OCH ₂ CF ₃	Me	H	Cl	Br	F	OCH ₂ CF ₃	Me	H	Cl
F	F	OCH ₂ CF ₃	Et	H	Cl	Br	F	OCH ₂ CF ₃	Et	H	Cl
F	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Br	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
F	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Br	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
F	F	OCH ₂ CF ₃	Me	Me	Cl	Br	F	OCH ₂ CF ₃	Me	Me	Cl
F	F	OCH ₂ CF ₃	Me	H	Br	Br	F	OCH ₂ CF ₃	Me	H	Br
F	F	OCH ₂ CF ₃	Et	H	Br	Br	F	OCH ₂ CF ₃	Et	H	Br
F	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Br	F	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
F	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Br	F	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
F	F	OCH ₂ CF ₃	Me	Me	Br	Br	F	OCH ₂ CF ₃	Me	Me	Br
F	Cl	CF ₃	Me	H	Cl	Br	Cl	CF ₃	Me	H	Cl
F	Cl	CF ₃	Et	H	Cl	Br	Cl	CF ₃	Et	H	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	Cl	CF ₃	<i>i</i> -Pr	H	Cl	Br	Cl	CF ₃	<i>i</i> -Pr	H	Cl
F	Cl	CF ₃	<i>t</i> -Bu	H	Cl	Br	Cl	CF ₃	<i>t</i> -Bu	H	Cl
F	Cl	CF ₃	Me	Me	Cl	Br	Cl	CF ₃	Me	Me	Cl
F	Cl	CF ₃	Me	H	Br	Br	Cl	CF ₃	Me	H	Br
F	Cl	CF ₃	Et	H	Br	Br	Cl	CF ₃	Et	H	Br
F	Cl	CF ₃	<i>i</i> -Pr	H	Br	Br	Cl	CF ₃	<i>i</i> -Pr	H	Br
F	Cl	CF ₃	<i>t</i> -Bu	H	Br	Br	Cl	CF ₃	<i>t</i> -Bu	H	Br
F	Cl	CF ₃	Me	Me	Br	Br	Cl	CF ₃	Me	Me	Br
F	Cl	Cl	Me	H	Cl	Br	Cl	Cl	Me	H	Cl
F	Cl	Cl	Et	H	Cl	Br	Cl	Cl	Et	H	Cl
F	Cl	Cl	<i>i</i> -Pr	H	Cl	Br	Cl	Cl	<i>i</i> -Pr	H	Cl
F	Cl	Cl	<i>t</i> -Bu	H	Cl	Br	Cl	Cl	<i>t</i> -Bu	H	Cl
F	Cl	Cl	Me	Me	Cl	Br	Cl	Cl	Me	Me	Cl
F	Cl	Cl	Me	H	Br	Br	Cl	Cl	Me	H	Br
F	Cl	Cl	Et	H	Br	Br	Cl	Cl	Et	H	Br
F	Cl	Cl	<i>i</i> -Pr	H	Br	Br	Cl	Cl	<i>i</i> -Pr	H	Br
F	Cl	Cl	<i>t</i> -Bu	H	Br	Br	Cl	Cl	<i>t</i> -Bu	H	Br
F	Cl	Cl	Me	Me	Br	Br	Cl	Cl	Me	Me	Br
F	Cl	Br	Me	H	Cl	Br	Cl	Br	Me	H	Cl
F	Cl	Br	Et	H	Cl	Br	Cl	Br	Et	H	Cl
F	Cl	Br	<i>i</i> -Pr	H	Cl	Br	Cl	Br	<i>i</i> -Pr	H	Cl
F	Cl	Br	<i>t</i> -Bu	H	Cl	Br	Cl	Br	<i>t</i> -Bu	H	Cl
F	Cl	Br	Me	Me	Cl	Br	Cl	Br	Me	Me	Cl
F	Cl	Br	Me	H	Br	Br	Cl	Br	Me	H	Br
F	Cl	Br	Et	H	Br	Br	Cl	Br	Et	H	Br
F	Cl	Br	<i>i</i> -Pr	H	Br	Br	Cl	Br	<i>i</i> -Pr	H	Br
F	Cl	Br	<i>t</i> -Bu	H	Br	Br	Cl	Br	<i>t</i> -Bu	H	Br
F	Cl	Br	Me	Me	Br	Br	Cl	Br	Me	Me	Br
F	Cl	OCH ₂ CF ₃	Me	H	Cl	Br	Cl	OCH ₂ CF ₃	Me	H	Cl
F	Cl	OCH ₂ CF ₃	Et	H	Cl	Br	Cl	OCH ₂ CF ₃	Et	H	Cl
F	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Br	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
F	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Br	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
F	Cl	OCH ₂ CF ₃	Me	Me	Cl	Br	Cl	OCH ₂ CF ₃	Me	Me	Cl
F	Cl	OCH ₂ CF ₃	Me	H	Br	Br	Cl	OCH ₂ CF ₃	Me	H	Br
F	Cl	OCH ₂ CF ₃	Et	H	Br	Br	Cl	OCH ₂ CF ₃	Et	H	Br
F	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Br	Cl	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
F	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Br	Cl	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	Cl	OCH ₂ CF ₃	Me	Me	Br	Br	Cl	OCH ₂ CF ₃	Me	Me	Br
F	Br	CF ₃	Me	H	Cl	Br	Br	CF ₃	Me	H	Cl
F	Br	CF ₃	Et	H	Cl	Br	Br	CF ₃	Et	H	Cl
F	Br	CF ₃	<i>i</i> -Pr	H	Cl	Br	Br	CF ₃	<i>i</i> -Pr	H	Cl
F	Br	CF ₃	<i>t</i> -Bu	H	Cl	Br	Br	CF ₃	<i>t</i> -Bu	H	Cl
F	Br	CF ₃	Me	Me	Cl	Br	Br	CF ₃	Me	Me	Cl
F	Br	CF ₃	Me	H	Br	Br	Br	CF ₃	Me	H	Br
F	Br	CF ₃	Et	H	Br	Br	Br	CF ₃	Et	H	Br
F	Br	CF ₃	<i>i</i> -Pr	H	Br	Br	Br	CF ₃	<i>i</i> -Pr	H	Br
F	Br	CF ₃	<i>t</i> -Bu	H	Br	Br	Br	CF ₃	<i>t</i> -Bu	H	Br
F	Br	CF ₃	Me	Me	Br	Br	Br	CF ₃	Me	Me	Br
F	Br	Cl	Me	H	Cl	Br	Br	Cl	Me	H	Cl
F	Br	Cl	Et	H	Cl	Br	Br	Cl	Et	H	Cl
F	Br	Cl	<i>i</i> -Pr	H	Cl	Br	Br	Cl	<i>i</i> -Pr	H	Cl
F	Br	Cl	<i>t</i> -Bu	H	Cl	Br	Br	Cl	<i>t</i> -Bu	H	Cl
F	Br	Cl	Me	Me	Cl	Br	Br	Cl	Me	Me	Cl
F	Br	Cl	Me	H	Br	Br	Br	Cl	Me	H	Br
F	Br	Cl	Et	H	Br	Br	Br	Cl	Et	H	Br
F	Br	Cl	<i>i</i> -Pr	H	Br	Br	Br	Cl	<i>i</i> -Pr	H	Br
F	Br	Cl	<i>t</i> -Bu	H	Br	Br	Br	Cl	<i>t</i> -Bu	H	Br
F	Br	Cl	Me	Me	Br	Br	Br	Cl	Me	Me	Br
F	Br	Br	Me	H	Cl	Br	Br	Br	Me	H	Cl
F	Br	Br	Et	H	Cl	Br	Br	Br	Et	H	Cl
F	Br	Br	<i>i</i> -Pr	H	Cl	Br	Br	Br	<i>i</i> -Pr	H	Cl
F	Br	Br	<i>t</i> -Bu	H	Cl	Br	Br	Br	<i>t</i> -Bu	H	Cl
F	Br	Br	Me	Me	Cl	Br	Br	Br	Me	Me	Cl
F	Br	Br	Me	H	Br	Br	Br	Br	Me	H	Br
F	Br	Br	Et	H	Br	Br	Br	Br	Et	H	Br
F	Br	Br	<i>i</i> -Pr	H	Br	Br	Br	Br	<i>i</i> -Pr	H	Br
F	Br	Br	<i>t</i> -Bu	H	Br	Br	Br	Br	<i>t</i> -Bu	H	Br
F	Br	Br	Me	Me	Br	Br	Br	Br	Me	Me	Br
F	Br	OCH ₂ CF ₃	Me	H	Cl	Br	Br	OCH ₂ CF ₃	Me	H	Cl
F	Br	OCH ₂ CF ₃	Et	H	Cl	Br	Br	OCH ₂ CF ₃	Et	H	Cl
F	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Br	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
F	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Br	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
F	Br	OCH ₂ CF ₃	Me	Me	Cl	Br	Br	OCH ₂ CF ₃	Me	Me	Cl
F	Br	OCH ₂ CF ₃	Me	H	Br	Br	Br	OCH ₂ CF ₃	Me	H	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	Br	OCH ₂ CF ₃	Et	H	Br	Br	Br	OCH ₂ CF ₃	Et	H	Br
F	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Br	Br	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
F	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Br	Br	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
F	Br	OCH ₂ CF ₃	Me	Me	Br	Br	Br	OCH ₂ CF ₃	Me	Me	Br
F	I	CF ₃	Me	H	Cl	Br	I	CF ₃	Me	H	Cl
F	I	CF ₃	Et	H	Cl	Br	I	CF ₃	Et	H	Cl
F	I	CF ₃	<i>i</i> -Pr	H	Cl	Br	I	CF ₃	<i>i</i> -Pr	H	Cl
F	I	CF ₃	<i>t</i> -Bu	H	Cl	Br	I	CF ₃	<i>t</i> -Bu	H	Cl
F	I	CF ₃	Me	Me	Cl	Br	I	CF ₃	Me	Me	Cl
F	I	CF ₃	Me	H	Br	Br	I	CF ₃	Me	H	Br
F	I	CF ₃	Et	H	Br	Br	I	CF ₃	Et	H	Br
F	I	CF ₃	<i>i</i> -Pr	H	Br	Br	I	CF ₃	<i>i</i> -Pr	H	Br
F	I	CF ₃	<i>t</i> -Bu	H	Br	Br	I	CF ₃	<i>t</i> -Bu	H	Br
F	I	CF ₃	Me	Me	Br	Br	I	CF ₃	Me	Me	Br
F	I	Cl	Me	H	Cl	Br	I	Cl	Me	H	Cl
F	I	Cl	Et	H	Cl	Br	I	Cl	Et	H	Cl
F	I	Cl	<i>i</i> -Pr	H	Cl	Br	I	Cl	<i>i</i> -Pr	H	Cl
F	I	Cl	<i>t</i> -Bu	H	Cl	Br	I	Cl	<i>t</i> -Bu	H	Cl
F	I	Cl	Me	Me	Cl	Br	I	Cl	Me	Me	Cl
F	I	Cl	Me	H	Br	Br	I	Cl	Me	H	Br
F	I	Cl	Et	H	Br	Br	I	Cl	Et	H	Br
F	I	Cl	<i>i</i> -Pr	H	Br	Br	I	Cl	<i>i</i> -Pr	H	Br
F	I	Cl	<i>t</i> -Bu	H	Br	Br	I	Cl	<i>t</i> -Bu	H	Br
F	I	Cl	Me	Me	Br	Br	I	Cl	Me	Me	Br
F	I	Br	Me	H	Cl	Br	I	Br	Me	H	Cl
F	I	Br	Et	H	Cl	Br	I	Br	Et	H	Cl
F	I	Br	<i>i</i> -Pr	H	Cl	Br	I	Br	<i>i</i> -Pr	H	Cl
F	I	Br	<i>t</i> -Bu	H	Cl	Br	I	Br	<i>t</i> -Bu	H	Cl
F	I	Br	Me	Me	Cl	Br	I	Br	Me	Me	Cl
F	I	Br	Me	H	Br	Br	I	Br	Me	H	Br
F	I	Br	Et	H	Br	Br	I	Br	Et	H	Br
F	I	Br	<i>i</i> -Pr	H	Br	Br	I	Br	<i>i</i> -Pr	H	Br
F	I	Br	<i>t</i> -Bu	H	Br	Br	I	Br	<i>t</i> -Bu	H	Br
F	I	Br	Me	Me	Br	Br	I	Br	Me	Me	Br
F	I	OCH ₂ CF ₃	Me	H	Cl	Br	I	OCH ₂ CF ₃	Me	H	Cl
F	I	OCH ₂ CF ₃	Et	H	Cl	Br	I	OCH ₂ CF ₃	Et	H	Cl
F	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Br	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Br	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
F	I	OCH ₂ CF ₃	Me	Me	Cl	Br	I	OCH ₂ CF ₃	Me	Me	Cl
F	I	OCH ₂ CF ₃	Me	H	Br	Br	I	OCH ₂ CF ₃	Me	H	Br
F	I	OCH ₂ CF ₃	Et	H	Br	Br	I	OCH ₂ CF ₃	Et	H	Br
F	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Br	I	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
F	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Br	I	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
F	I	OCH ₂ CF ₃	Me	Me	Br	Br	I	OCH ₂ CF ₃	Me	Me	Br
F	CF ₃	CF ₃	Me	H	Cl	Br	CF ₃	CF ₃	Me	H	Cl
F	CF ₃	CF ₃	Et	H	Cl	Br	CF ₃	CF ₃	Et	H	Cl
F	CF ₃	CF ₃	<i>i</i> -Pr	H	Cl	Br	CF ₃	CF ₃	<i>i</i> -Pr	H	Cl
F	CF ₃	CF ₃	<i>t</i> -Bu	H	Cl	Br	CF ₃	CF ₃	<i>t</i> -Bu	H	Cl
F	CF ₃	CF ₃	Me	Me	Cl	Br	CF ₃	CF ₃	Me	Me	Cl
F	CF ₃	CF ₃	Me	H	Br	Br	CF ₃	CF ₃	Me	H	Br
F	CF ₃	CF ₃	Et	H	Br	Br	CF ₃	CF ₃	Et	H	Br
F	CF ₃	CF ₃	<i>i</i> -Pr	H	Br	Br	CF ₃	CF ₃	<i>i</i> -Pr	H	Br
F	CF ₃	CF ₃	<i>t</i> -Bu	H	Br	Br	CF ₃	CF ₃	<i>t</i> -Bu	H	Br
F	CF ₃	CF ₃	Me	Me	Br	Br	CF ₃	CF ₃	Me	Me	Br
F	CF ₃	Cl	Me	H	Cl	Br	CF ₃	Cl	Me	H	Cl
F	CF ₃	Cl	Et	H	Cl	Br	CF ₃	Cl	Et	H	Cl
F	CF ₃	Cl	<i>i</i> -Pr	H	Cl	Br	CF ₃	Cl	<i>i</i> -Pr	H	Cl
F	CF ₃	Cl	<i>t</i> -Bu	H	Cl	Br	CF ₃	Cl	<i>t</i> -Bu	H	Cl
F	CF ₃	Cl	Me	Me	Cl	Br	CF ₃	Cl	Me	Me	Cl
F	CF ₃	Cl	Me	H	Br	Br	CF ₃	Cl	Me	H	Br
F	CF ₃	Cl	Et	H	Br	Br	CF ₃	Cl	Et	H	Br
F	CF ₃	Cl	<i>i</i> -Pr	H	Br	Br	CF ₃	Cl	<i>i</i> -Pr	H	Br
F	CF ₃	Cl	<i>t</i> -Bu	H	Br	Br	CF ₃	Cl	<i>t</i> -Bu	H	Br
F	CF ₃	Cl	Me	Me	Br	Br	CF ₃	Cl	Me	Me	Br
F	CF ₃	Br	Me	H	Cl	Br	CF ₃	Br	Me	H	Cl
F	CF ₃	Br	Et	H	Cl	Br	CF ₃	Br	Et	H	Cl
F	CF ₃	Br	<i>i</i> -Pr	H	Cl	Br	CF ₃	Br	<i>i</i> -Pr	H	Cl
F	CF ₃	Br	<i>t</i> -Bu	H	Cl	Br	CF ₃	Br	<i>t</i> -Bu	H	Cl
F	CF ₃	Br	Me	Me	Cl	Br	CF ₃	Br	Me	Me	Cl
F	CF ₃	Br	Me	H	Br	Br	CF ₃	Br	Me	H	Br
F	CF ₃	Br	Et	H	Br	Br	CF ₃	Br	Et	H	Br
F	CF ₃	Br	<i>i</i> -Pr	H	Br	Br	CF ₃	Br	<i>i</i> -Pr	H	Br
F	CF ₃	Br	<i>t</i> -Bu	H	Br	Br	CF ₃	Br	<i>t</i> -Bu	H	Br
F	CF ₃	Br	Me	Me	Br	Br	CF ₃	Br	Me	Me	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	CF ₃	OCH ₂ CF ₃	Me	H	Cl	Br	CF ₃	OCH ₂ CF ₃	Me	H	Cl
F	CF ₃	OCH ₂ CF ₃	Et	H	Cl	Br	CF ₃	OCH ₂ CF ₃	Et	H	Cl
F	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl	Br	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Cl
F	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl	Br	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Cl
F	CF ₃	OCH ₂ CF ₃	Me	Me	Cl	Br	CF ₃	OCH ₂ CF ₃	Me	Me	Cl
F	CF ₃	OCH ₂ CF ₃	Me	H	Br	Br	CF ₃	OCH ₂ CF ₃	Me	H	Br
F	CF ₃	OCH ₂ CF ₃	Et	H	Br	Br	CF ₃	OCH ₂ CF ₃	Et	H	Br
F	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br	Br	CF ₃	OCH ₂ CF ₃	<i>i</i> -Pr	H	Br
F	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br	Br	CF ₃	OCH ₂ CF ₃	<i>t</i> -Bu	H	Br
F	CF ₃	OCH ₂ CF ₃	Me	Me	Br	Br	CF ₃	OCH ₂ CF ₃	Me	Me	Br
CH ₃	F	OCHF ₂	Me	H	Cl	Cl	F	OCHF ₂	Me	H	Cl
CH ₃	F	OCHF ₂	Et	H	Cl	Cl	F	OCHF ₂	Et	H	Cl
CH ₃	F	OCHF ₂	<i>i</i> -Pr	H	Cl	Cl	F	OCHF ₂	<i>i</i> -Pr	H	Cl
CH ₃	F	OCHF ₂	<i>t</i> -Bu	H	Cl	Cl	F	OCHF ₂	<i>t</i> -Bu	H	Cl
CH ₃	F	OCHF ₂	Me	Me	Cl	Cl	F	OCHF ₂	Me	Me	Cl
CH ₃	F	OCHF ₂	Me	H	Br	Cl	F	OCHF ₂	Me	H	Br
CH ₃	F	OCHF ₂	Et	H	Br	Cl	F	OCHF ₂	Et	H	Br
CH ₃	F	OCHF ₂	<i>i</i> -Pr	H	Br	Cl	F	OCHF ₂	<i>i</i> -Pr	H	Br
CH ₃	F	OCHF ₂	<i>t</i> -Bu	H	Br	Cl	F	OCHF ₂	<i>t</i> -Bu	H	Br
CH ₃	F	OCHF ₂	Me	Me	Br	Cl	F	OCHF ₂	Me	Me	Br
CH ₃	Cl	OCHF ₂	Me	H	Cl	Cl	Cl	OCHF ₂	Me	H	Cl
CH ₃	Cl	OCHF ₂	Et	H	Cl	Cl	Cl	OCHF ₂	Et	H	Cl
CH ₃	Cl	OCHF ₂	<i>i</i> -Pr	H	Cl	Cl	Cl	OCHF ₂	<i>i</i> -Pr	H	Cl
CH ₃	Cl	OCHF ₂	<i>t</i> -Bu	H	Cl	Cl	Cl	OCHF ₂	<i>t</i> -Bu	H	Cl
CH ₃	Cl	OCHF ₂	Me	Me	Cl	Cl	Cl	OCHF ₂	Me	Me	Cl
CH ₃	Cl	OCHF ₂	Me	H	Br	Cl	Cl	OCHF ₂	Me	H	Br
CH ₃	Cl	OCHF ₂	Et	H	Br	Cl	Cl	OCHF ₂	Et	H	Br
CH ₃	Cl	OCHF ₂	<i>i</i> -Pr	H	Br	Cl	Cl	OCHF ₂	<i>i</i> -Pr	H	Br
CH ₃	Cl	OCHF ₂	<i>t</i> -Bu	H	Br	Cl	Cl	OCHF ₂	<i>t</i> -Bu	H	Br
CH ₃	Cl	OCHF ₂	Me	Me	Br	Cl	Cl	OCHF ₂	Me	Me	Br
CH ₃	Br	OCHF ₂	Me	H	Cl	Cl	Br	OCHF ₂	Me	H	Cl
CH ₃	Br	OCHF ₂	Et	H	Cl	Cl	Br	OCHF ₂	Et	H	Cl
CH ₃	Br	OCHF ₂	<i>i</i> -Pr	H	Cl	Cl	Br	OCHF ₂	<i>i</i> -Pr	H	Cl
CH ₃	Br	OCHF ₂	<i>t</i> -Bu	H	Cl	Cl	Br	OCHF ₂	<i>t</i> -Bu	H	Cl
CH ₃	Br	OCHF ₂	Me	Me	Cl	Cl	Br	OCHF ₂	Me	Me	Cl
CH ₃	Br	OCHF ₂	Me	H	Br	Cl	Br	OCHF ₂	Me	H	Br
CH ₃	Br	OCHF ₂	Et	H	Br	Cl	Br	OCHF ₂	Et	H	Br

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
CH ₃	Br	OCHF ₂	<i>i</i> -Pr	H	Br	Cl	Br	OCHF ₂	<i>i</i> -Pr	H	Br
CH ₃	Br	OCHF ₂	<i>t</i> -Bu	H	Br	Cl	Br	OCHF ₂	<i>t</i> -Bu	H	Br
CH ₃	Br	OCHF ₂	Me	Me	Br	Cl	Br	OCHF ₂	Me	Me	Br
CH ₃	I	OCHF ₂	Me	H	Cl	Cl	I	OCHF ₂	Me	H	Cl
CH ₃	I	OCHF ₂	Et	H	Cl	Cl	I	OCHF ₂	Et	H	Cl
CH ₃	I	OCHF ₂	<i>i</i> -Pr	H	Cl	Cl	I	OCHF ₂	<i>i</i> -Pr	H	Cl
CH ₃	I	OCHF ₂	<i>t</i> -Bu	H	Cl	Cl	I	OCHF ₂	<i>t</i> -Bu	H	Cl
CH ₃	I	OCHF ₂	Me	Me	Cl	Cl	I	OCHF ₂	Me	Me	Cl
CH ₃	I	OCHF ₂	Me	H	Br	Cl	I	OCHF ₂	Me	H	Br
CH ₃	I	OCHF ₂	Et	H	Br	Cl	I	OCHF ₂	Et	H	Br
CH ₃	I	OCHF ₂	<i>i</i> -Pr	H	Br	Cl	I	OCHF ₂	<i>i</i> -Pr	H	Br
CH ₃	I	OCHF ₂	<i>t</i> -Bu	H	Br	Cl	I	OCHF ₂	<i>t</i> -Bu	H	Br
CH ₃	I	OCHF ₂	Me	Me	Br	Cl	I	OCHF ₂	Me	Me	Br
CH ₃	CF ₃	OCHF ₂	Me	H	Cl	Cl	CF ₃	OCHF ₂	Me	H	Cl
CH ₃	CF ₃	OCHF ₂	Et	H	Cl	Cl	CF ₃	OCHF ₂	Et	H	Cl
CH ₃	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Cl	Cl	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Cl
CH ₃	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Cl	Cl	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Cl
CH ₃	CF ₃	OCHF ₂	Me	Me	Cl	Cl	CF ₃	OCHF ₂	Me	Me	Cl
CH ₃	CF ₃	OCHF ₂	Me	H	Br	Cl	CF ₃	OCHF ₂	Me	H	Br
CH ₃	CF ₃	OCHF ₂	Et	H	Br	Cl	CF ₃	OCHF ₂	Et	H	Br
CH ₃	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Br	Cl	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Br
CH ₃	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Br	Cl	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Br
CH ₃	CF ₃	OCHF ₂	Me	Me	Br	Cl	CF ₃	OCHF ₂	Me	Me	Br
F	F	OCHF ₂	Me	H	Cl	Br	F	OCHF ₂	Me	H	Cl
F	F	OCHF ₂	Et	H	Cl	Br	F	OCHF ₂	Et	H	Cl
F	F	OCHF ₂	<i>i</i> -Pr	H	Cl	Br	F	OCHF ₂	<i>i</i> -Pr	H	Cl
F	F	OCHF ₂	<i>t</i> -Bu	H	Cl	Br	F	OCHF ₂	<i>t</i> -Bu	H	Cl
F	F	OCHF ₂	Me	Me	Cl	Br	F	OCHF ₂	Me	Me	Cl
F	F	OCHF ₂	Me	H	Br	Br	F	OCHF ₂	Me	H	Br
F	F	OCHF ₂	Et	H	Br	Br	F	OCHF ₂	Et	H	Br
F	F	OCHF ₂	<i>i</i> -Pr	H	Br	Br	F	OCHF ₂	<i>i</i> -Pr	H	Br
F	F	OCHF ₂	<i>t</i> -Bu	H	Br	Br	F	OCHF ₂	<i>t</i> -Bu	H	Br
F	F	OCHF ₂	Me	Me	Br	Br	F	OCHF ₂	Me	Me	Br
F	Cl	OCHF ₂	Me	H	Cl	Br	Cl	OCHF ₂	Me	H	Cl
F	Cl	OCHF ₂	Et	H	Cl	Br	Cl	OCHF ₂	Et	H	Cl
F	Cl	OCHF ₂	<i>i</i> -Pr	H	Cl	Br	Cl	OCHF ₂	<i>i</i> -Pr	H	Cl
F	Cl	OCHF ₂	<i>t</i> -Bu	H	Cl	Br	Cl	OCHF ₂	<i>t</i> -Bu	H	Cl

<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R⁹</u>	<u>R³</u>	<u>R²</u>	<u>R⁷</u>
F	Cl	OCHF ₂	Me	Me	Cl	Br	Cl	OCHF ₂	Me	Me	Cl
F	Cl	OCHF ₂	Me	H	Br	Br	Cl	OCHF ₂	Me	H	Br
F	Cl	OCHF ₂	Et	H	Br	Br	Cl	OCHF ₂	Et	H	Br
F	Cl	OCHF ₂	<i>i</i> -Pr	H	Br	Br	Cl	OCHF ₂	<i>i</i> -Pr	H	Br
F	Cl	OCHF ₂	<i>t</i> -Bu	H	Br	Br	Cl	OCHF ₂	<i>t</i> -Bu	H	Br
F	Cl	OCHF ₂	Me	Me	Br	Br	Cl	OCHF ₂	Me	Me	Br
F	Br	OCHF ₂	Me	H	Cl	Br	Br	OCHF ₂	Me	H	Cl
F	Br	OCHF ₂	Et	H	Cl	Br	Br	OCHF ₂	Et	H	Cl
F	Br	OCHF ₂	<i>i</i> -Pr	H	Cl	Br	Br	OCHF ₂	<i>i</i> -Pr	H	Cl
F	Br	OCHF ₂	<i>t</i> -Bu	H	Cl	Br	Br	OCHF ₂	<i>t</i> -Bu	H	Cl
F	Br	OCHF ₂	Me	Me	Cl	Br	Br	OCHF ₂	Me	Me	Cl
F	Br	OCHF ₂	Me	H	Br	Br	Br	OCHF ₂	Me	H	Br
F	Br	OCHF ₂	Et	H	Br	Br	Br	OCHF ₂	Et	H	Br
F	Br	OCHF ₂	<i>i</i> -Pr	H	Br	Br	Br	OCHF ₂	<i>i</i> -Pr	H	Br
F	Br	OCHF ₂	<i>t</i> -Bu	H	Br	Br	Br	OCHF ₂	<i>t</i> -Bu	H	Br
F	Br	OCHF ₂	Me	Me	Br	Br	Br	OCHF ₂	Me	Me	Br
F	I	OCHF ₂	Me	H	Cl	Br	I	OCHF ₂	Me	H	Cl
F	I	OCHF ₂	Et	H	Cl	Br	I	OCHF ₂	Et	H	Cl
F	I	OCHF ₂	<i>i</i> -Pr	H	Cl	Br	I	OCHF ₂	<i>i</i> -Pr	H	Cl
F	I	OCHF ₂	<i>t</i> -Bu	H	Cl	Br	I	OCHF ₂	<i>t</i> -Bu	H	Cl
F	I	OCHF ₂	Me	Me	Cl	Br	I	OCHF ₂	Me	Me	Cl
F	I	OCHF ₂	Me	H	Br	Br	I	OCHF ₂	Me	H	Br
F	I	OCHF ₂	Et	H	Br	Br	I	OCHF ₂	Et	H	Br
F	I	OCHF ₂	<i>i</i> -Pr	H	Br	Br	I	OCHF ₂	<i>i</i> -Pr	H	Br
F	I	OCHF ₂	<i>t</i> -Bu	H	Br	Br	I	OCHF ₂	<i>t</i> -Bu	H	Br
F	I	OCHF ₂	Me	Me	Br	Br	I	OCHF ₂	Me	Me	Br
F	CF ₃	OCHF ₂	Me	H	Cl	Br	CF ₃	OCHF ₂	Me	H	Cl
F	CF ₃	OCHF ₂	Et	H	Cl	Br	CF ₃	OCHF ₂	Et	H	Cl
F	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Cl	Br	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Cl
F	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Cl	Br	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Cl
F	CF ₃	OCHF ₂	Me	Me	Cl	Br	CF ₃	OCHF ₂	Me	Me	Cl
F	CF ₃	OCHF ₂	Me	H	Br	Br	CF ₃	OCHF ₂	Me	H	Br
F	CF ₃	OCHF ₂	Et	H	Br	Br	CF ₃	OCHF ₂	Et	H	Br
F	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Br	Br	CF ₃	OCHF ₂	<i>i</i> -Pr	H	Br
F	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Br	Br	CF ₃	OCHF ₂	<i>t</i> -Bu	H	Br
F	CF ₃	OCHF ₂	Me	Me	Br	Br	CF ₃	OCHF ₂	Me	Me	Br

CLAIMS

What is claimed is:


1. A method for preparing a fused oxazinone, comprising:

5 (1) contacting a carboxylic acid with a sulfonyl chloride in the presence of an optionally substituted pyridine compound, the nominal mole ratio of sulfonyl chloride to carboxylic acid being from about 0.75 to 1.5;

10 (2) contacting the mixture prepared in (1) with an *ortho*-amino aromatic carboxylic acid in the presence of an optionally substituted pyridine compound, the nominal mole ratio of the *ortho*-amino aromatic carboxylic acid to carboxylic acid charged in (1) being from about 0.8 to 1.2; and

(3) adding additional sulfonyl chloride to the mixture prepared in (2), the nominal mole ratio of additional sulfonyl chloride added in (3) to carboxylic acid charged in (1) being at least about 0.5.

2. The method of Claim 1 wherein a fused oxazinone of Formula 1



15


wherein

J is an optionally substituted carbon moiety; and


K is, together with the two contiguous linking carbon atoms, a fused phenyl ring or a fused 5- or 6-membered heteroaromatic ring, each ring optionally substituted;

20 is prepared by

(1) contacting a carboxylic acid of Formula 2




with a sulfonyl chloride of Formula 4



25 wherein L is selected from alkyl, haloalkyl, and phenyl optionally substituted with from one to three substituents independently selected from alkyl or halogen;

in the presence of an optionally substituted pyridine compound;

(2) contacting the mixture prepared in (1) with an *ortho*-amino carboxylic acid of Formula 5



in the presence of an optionally substituted pyridine compound; and
(3) contacting the mixture prepared in (2) with additional sulfonyl chloride of Formula 4.

3. The method of Claim 2 wherein the nominal mole ratio of sulfonyl chloride to
5 carboxylic acid in (1) is from about 1.0 to 1.5; the nominal mole ratio of the *ortho*-amino
aromatic carboxylic acid in (2) to carboxylic acid charged in (1) is from about 0.9 to 1.1; the
nominal mole ratio of additional sulfonyl chloride added in (3) to carboxylic acid charged in
(1) is from about 1.0 to 1.5.

4. The method of Claim 3 wherein the nominal mole ratio of the optionally
10 substituted pyridine compound charged in (1) to carboxylic acid charged in (1) is from about
1.0 to 2.0; additional optionally substituted pyridine compound is charged in (2); and the
nominal mole ratio of the additional optionally substituted pyridine compound charged in
(2) to carboxylic acid charged in (1) is from about 2.0 to 4.0.

5. The method of Claim 2 wherein
15 J is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₈
cycloalkenyl, each optionally substituted; or
J is a phenyl ring, a benzyl group, a benzoyl group, a 5- or 6-membered heteroaromatic
ring, an aromatic 8-, 9- or 10-membered fused carbobicyclic ring system, an
aromatic 8-, 9- or 10-membered fused heterobicyclic ring system or a 5- or 6-
20 membered nonaromatic heterocyclic ring optionally including one or two ring
members selected from the group consisting of C(=O), SO or S(O)₂, each
optionally substituted.

6. The method of Claim 5 wherein
K is, together with the two contiguous linking carbon atoms, a fused phenyl ring
25 optionally substituted with from one to four substituents independently selected
from G, U, W or R¹³; or a fused 5- or 6-membered heteroaromatic ring
optionally substituted with from one to three substituents independently selected
from G, U, W or R¹³;
J is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₈
30 cycloalkenyl, each optionally substituted with one or more substituents selected
from the group consisting of R¹², halogen, CN, NO₂, hydroxy, C₁-C₄ alkoxy,
C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈

dialkylamino, C₃-C₆ cycloalkylamino, and (C₁-C₄ alkyl)C₃-C₆ cycloalkylamino;
or

- J is a phenyl ring, a benzyl group, a benzoyl group, a 5- or 6-membered heteroaromatic ring, an aromatic 8-, 9- or 10-membered fused carbobicyclic ring system, an aromatic 8-, 9- or 10-membered fused heterobicyclic ring system or a 5- or 6-membered nonaromatic heterocyclic ring optionally including one or two ring members selected from the group consisting of C(=O), SO or S(O)₂, each optionally substituted with from one to four substituents independently selected from G, U, W or R¹³;
- each G is a 5- or 6-membered nonaromatic heterocyclic ring optionally including one or two ring members selected from the group consisting of C(=O), SO or S(O)₂, each optionally substituted with from one to four substituents independently selected from W;
- each U is a phenyl ring, a benzyl group, a benzoyl group, a 5- or 6-membered heteroaromatic ring, an aromatic 8-, 9- or 10-membered fused carbobicyclic ring system, an aromatic 8-, 9- or 10-membered fused heterobicyclic ring system, each optionally substituted with from one to four substituents independently selected from W;
- each W is C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, C₃-C₆ cycloalkylamino, (C₁-C₄ alkyl)C₃-C₆ cycloalkylamino and C₃-C₆ trialkylsilyl;
- each R¹² is independently R¹⁹C(=E)-; R¹⁹C(=E)L-; R¹⁹LC(=E)-; (R¹⁹)LC(=E)L-; -O(Q=)P(OR¹⁹)₂; -SO₂LR¹⁸; or R¹⁹SO₂L-;
- each R¹³ is B(OR¹⁷)₂; NH₂; SH; thiocyanato; C₃-C₈ trialkylsilyloxy; C₁-C₄ alkyldisulfide; SF₅; R¹⁹C(=E)-; R¹⁹C(=E)M-; R¹⁹MC(=E)-; (R¹⁹)MC(=E)M-; -OP(=Q)(OR¹⁹)₂; -SO₂MR¹⁹; R¹⁹SO₂M-;
- each E is independently O, S, NR¹⁵, NOR¹⁵, NN(R¹⁵)₂, N-S=O, N-CN or N-NO₂;
- each M is independently O, NR¹⁸ or S;
- Q is O or S;
- each R¹⁵ and each R¹⁹ is independently H; C₁-C₆ alkyl optionally substituted with one or more substituents selected from the group consisting of CN, NO₂, hydroxy, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkylthio, C₁-C₄ haloalkylsulfinyl, C₁-C₄ haloalkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino, CO₂H, C₂-C₆ alkoxycarbonyl, C₂-C₆ alkylcarbonyl, C₃-C₆ trialkylsilyl, and a phenyl ring


optionally substituted with one to three substituents independently selected from W; C₁-C₆ haloalkyl; C₃-C₆ cycloalkyl; or a phenyl ring optionally substituted with from one to three substituents independently selected from W;

each R¹⁷ is independently H or C₁-C₄ alkyl; or

- 5 B(OR¹⁷)₂ can form a ring wherein the two oxygen atoms are linked by a chain of two to three carbons optionally substituted with one or two substituents independently selected from methyl or C₂-C₆ alkoxy carbonyl; and each R¹⁸ is independently H, C₁-C₆ alkyl or C₁-C₆ haloalkyl.

7. The method of Claim 6 wherein K is, together with the two contiguous linking carbon atoms, a fused phenyl ring optionally substituted with from one to four substituents independently selected from W or R¹³.


8. The method of Claim 2 wherein a compound of Formula 1a



wherein

- 15 X is N or CR⁶;
Y is N or CH;
R⁴ is C₁-C₄ alkyl or halogen;
R⁵ is H, C₁-C₄ alkyl, C₁-C₄ haloalkyl or halogen;
R⁶ and R⁷ are independently H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, halogen, CN or C₁-C₄
20 haloalkoxy;
R⁸ is H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ halocycloalkyl, halogen, CN, NO₂, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylamino, C₂-C₈ dialkylamino,
25 C₃-C₆ cycloalkylamino, (C₁-C₄ alkyl)(C₃-C₆ cycloalkyl)amino, C₂-C₄ alkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₂-C₆ alkylaminocarbonyl, C₃-C₈ dialkylaminocarbonyl or C₃-C₆ trialkylsilyl; and
R⁹ is CF₃, OCF₃, OCHF₂, OCH₂CF₃, S(O)_pCF₃, S(O)_pCHF₂ or halogen;

is prepared using a compound of Formula 2' as the Formula 2 compound and a compound of Formula 5' as the Formula 5 compound



9. The method of Claim 8 wherein

X is N;

Y is N;

R⁴ is CH₃, F, Cl or Br;


R⁵ is CF₃, F, Cl, Br or I;

R⁷ is Cl or Br;

R⁸ is H, and

R⁹ is CF₃, OCHF₂, OCH₂CF₃, Cl or Br.

10. A method for preparing a compound of Formula III.



wherein

X is N or CR⁶;

Y is N or CH;

R¹ is H;

R² is H or CH₃;

R³ is C₁-C₆ alkyl;


R⁴ is C₁-C₄ alkyl or halogen;

R⁵ is H, C₁-C₄ alkyl, C₁-C₄ haloalkyl or halogen;

R^6 and R^7 are independently H, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen, CN or C_1 - C_4 haloalkoxy;

R^8 is H, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_6 cycloalkyl, C_1 - C_4 haloalkyl, C_2 - C_4 haloalkenyl, C_2 - C_4 haloalkynyl, C_3 - C_6 halocycloalkyl, halogen, CN, NO_2 , C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 alkylamino, C_2 - C_8 dialkylamino, C_3 - C_6 cycloalkylamino, $(C_1$ - C_4 alkyl)(C_3 - C_6 cycloalkyl)amino, C_2 - C_4 alkylcarbonyl, C_2 - C_6 alkoxy carbonyl, C_2 - C_6 alkylaminocarbonyl, C_3 - C_8 dialkylaminocarbonyl or C_3 - C_6 trialkylsilyl; and

R^9 is CF_3 , OCF_3 , $OCHF_2$, OCH_2CF_3 , $S(O)_pCF_3$, $S(O)_pCHF_2$ or halogen; using a compound of Formula 1a



characterized by:

preparing said compound of Formula 1a by the method of Claim 8.

11. The method of Claim 10 wherein

X is N;

Y is N;

R^2 is H or CH_3 ;

R^3 is C_1 - C_4 alkyl;

R^4 is CH_3 , F, Cl or Br;

R^5 is CF_3 , F, Cl, Br or I;

R^7 is Cl or Br;

R^8 is H, and

R^9 is CF_3 , $OCHF_2$, OCH_2CF_3 , Cl or Br.

TITLE

METHOD FOR PREPARING FUSED OXAZINONES

ABSTRACT OF THE DISCLOSURE


A method for preparing a fused oxazinone is disclosed in which

5 (1) a carboxylic acid is contacted with a sulfonyl chloride in the presence of an optionally substituted pyridine compound, the nominal mole ratio of sulfonyl chloride to carboxylic acid being from about 0.75 to 1.5;

(2) the mixture prepared in (1) is contacted with an *ortho*-amino aromatic carboxylic acid in the presence of an optionally substituted pyridine compound, the nominal mole ratio
10 of the *ortho*-amino aromatic carboxylic acid to carboxylic acid charged in (1) being from about 0.8 to 1.2; and

(3) additional sulfonyl chloride is added to the mixture prepared in (2), the nominal mole ratio of additional sulfonyl chloride added in (3) to carboxylic acid charged in (1) being at least about 0.5.

15 Also disclosed is a method for preparing a compound of Formula III, using a compound of Formula 1a that is characterized by preparing the fused oxazinone of Formula 1a by the method above, using a compound of the formula $LS(O)_2Cl$ as the sulfonyl chloride, a compound of Formula 2' as the carboxylic acid, and a compound of Formula 5' as the *ortho*-amino aromatic carboxylic acid



wherein L, X, Y and R¹ through R⁹ are as defined in the disclosure.